

```
$ EOLCOM >
$ TITLE RIO GRANDE BASIN HYDROECONOMIC PROTOTYPE
$ OFFSYMREF OFFSYMLIST OFFLISTING OFFUPPER
```

```
OPTION LIMROW = 0, LIMCOL = 0;
OPTION iterlim = 10000000;
```

```
$ONTEXT
```

```
* -----
Output control commands above vary listing's appearance
```

```
EOLCOM > tells GAMS to ignore anything in the line's text after >
OFFLISTING deletes all program lines and just includes GAMS listing
Setting LIMROW = 0 eliminates all equations in the GAMS listing
It saves space, but is usually a bad idea till the model is known bullet proof
```

```
Colors: We suggest going to 'file' then to 'options,'
then choose as many colors as possible for varying kinds of GAMS syntax
It simplifies error trapping.
```

```
* -----
August 29, 2006
```

```
Rio Grande Basin Model with crops:
```

```
Contains most elements of full Upper Rio Grande Basin Model from 2001-2006.
Complete hydrological mass balance for all surface water and groundwater
All use, through diversions or pumping, deplete the river and/or its connected aquifer
All economic benefits derive from agriculture, M&I, and reservoir recreation
Endangered species needs set minimum flows
Terminal conditions are specified for flow, aquifers, and reservoirs
```

```
Crop and irrigation technology choices are also included in this model
```

```
* -----
Originally developed by Dr. Jim Booker (1996-2001), and funded by
```

```
US Geological Survey,
Water Resources Research Institutes and Ag Experiment Stations of CO, NM, and TX
```

```
Currently (2001-2006) sponsored by Rio Grande Basin Initiative (USDA and US Congress)
```

```
Model Contacts: Frank Ward, New Mexico State University: fward@nmsu.edu
Jim Booker, Siena College, New York: jbooker@siena.edu
Hilary Brinegar, New Mexico State Univ: hilarybrinegar@hotmail.com
```

```
* -----
Model has these flow nodes:
```

```
9 inflow headwater nodes
11 river gauge nodes
7 diversion nodes
7 application nodes
7 consumptive use nodes
```

7 seepage nodes  
7 net seepage nodes  
7 pumping nodes  
7 surface water to river return flow nodes

7 groundwater to river return flow nodes  
7 groundwater to aquifer volume gain nodes

6 reservoir release to river node  
6 reservoir evaporation nodes

and these stock nodes:

6 reservoir node  
7 aquifer nodes

\* -----

*FLOWS: Spatial unit for FLOWS is set (index) i.*

*Each element in the set i is assigned to one water use subset (category)*

*Subset categories include:*

- |   |                    |
|---|--------------------|
| 1. Inflow (headwater) nodes to the system,                | <i>inflow(i);</i>  |
| 2. Nodes on a river or tributary                          | <i>river(i);</i>   |
| 3. Diversion nodes  | <i>divert(i);</i>  |
| 4. Application nodes                                      | <i>apply(i);</i>   |
| 5. Consumptive use nodes                                  | <i>use(i);</i>     |
| 6. Seepage nodes  | <i>seep(i);</i>    |
| 7. Net Seepage nodes                                      | <i>netseep(i);</i> |
| 8. Pumping nodes  | <i>pump(i);</i>    |
| 9. Surface return flow nodes directly to the river,       | <i>return(i);</i>  |
| 10. Groundwater return flow groundwater net seepage       | <i>gwflow(i);</i>  |
| 11. Gain in aquifer volume nodes from net seepage         | <i>aqflow(i);</i>  |
| 12. NET reservoir releases from storage, outflow - inflow | <i>rel(i);</i>     |
| 13. reservoir evaporation nodes, based on surf area       | <i>evp(i);</i>     |

*STOCKS: Spatial unit for STOCKS is the set index u.*

*Each element of the set u is assigned to one water use subset (category).*

*Subset categories are:*

- |                           |                |
|---------------------------|----------------|
| 1. Reservoir volume nodes | <i>res(u);</i> |
| 2. Aquifer volume node    | <i>aqf(u);</i> |

\* -----

TABLE OF CONTENTS

- Section 1. Sets
- Section 2. Data
- Section 3. Variables
- Section 4. Equations
- Section 5. Models
- Section 6. Solves
- Section 7. Displays

\* -----

\$OFFTEXT

\*\*\*\*\* Section 1 \*\*\*\*\*  
\* The following sets are specified as indices \*  
\* for parameters (data), variables, and equations \*  
\*\*\*\*\*

SETS

\*\*\*\*\*  
i Flows -- location of important nodes in RG Basin -- Colorado to Mexico  
\*\*\*\*\*

/	RG-DN_h_f	Headwater flow nodes	inflow(i)
	Conejos_h_f		
	CBasn_h_f		
	SangDC_h_f		
	SJCham_h_f		
	Chama_h_f		
	Jemez_h_f		
	Puerco_h_f		
	Salado_h_f		
	Lobatos_v_f	River gage measurement nodes	river(i)
	Embudo_v_f		
	Chamita_v_f		
	Otowi_v_f		
	Coch_g_v_f		
	Acacia_v_f		
	Marcial_v_f		
	EB_g_v_f		
	CA_g_v_f		
	ElPaso_v_f		
	Quitman_v_f		
	SLV_f	All agriculture nodes	ag(i)
	MRGCD_f		
	EBID_f		
	MXAG_f		
	EPAG_f		
	ABQMI_f	All M&I nodes	mi(i)
	EPMI_f		
	SLV_d_f	Diversion nodes	divert(i)

ABQMI\_d\_f  
MRGCD\_d\_f  
EBID\_d\_f  
MXAG\_d\_f  
EPMI\_d\_f  
EPAG\_d\_f

*Application nodes*

*apply(i)*

SLV\_a\_f  
ABQMI\_a\_f  
MRGCD\_a\_f  
EBID\_a\_f  
MXAG\_a\_f  
EPMI\_a\_f  
EPAG\_a\_f

*Consumptive use (ET) nodes*

*use(i)*

SLV\_u\_f  
ABQMI\_u\_f  
MRGCD\_u\_f  
EBID\_u\_f  
MXAG\_u\_f  
EPMI\_u\_f  
EPAG\_u\_f

*Seepage nodes*

*seep(i)*

SLV\_s\_f  
ABQMI\_s\_f  
MRGCD\_s\_f  
EBID\_s\_f  
MXAG\_s\_f  
EPMI\_s\_f  
EPAG\_s\_f

*Pumping nodes*

*pump(i)*

SLV\_p\_f  
ABQMI\_p\_f  
MRGCD\_p\_f  
EBID\_p\_f  
MXAG\_p\_f  
EPMI\_p\_f  
EPAG\_p\_f

*Net Seepage nodes*

*netseep(i)*

SLV\_n\_f  
ABQMI\_n\_f  
MRGCD\_n\_f  
EBID\_n\_f  
MXAG\_n\_f  
EPMI\_n\_f  
EPAG\_n\_f

*Return flow (surface) nodes*

*return(i)*

SLV\_r\_f  
ABQMI\_r\_f  
MRGCD\_r\_f  
EBID\_r\_f  
MXAG\_r\_f  
EPMI\_r\_f  
EPAG\_r\_f

*Groundwater return flow to riv nodes*

*gwflow(i)*

SLV\_g\_f

```

ABQMI_g_f
MRGCD_g_f
EBID_g_f
MXAG_g_f
EPMI_g_f
EPAG_g_f

SLV_dq_f      Gain in aquifer volume nodes      aqflow(i)
ABQMI_dq_f
MRGCD_dq_f
EBID_dq_f
MXAG_dq_f
EPMI_dq_f
EPAG_dq_f

HE_rel_f      Reservoir-to-river release flow nodes  release(i)
EV_rel_f
AB_rel_f
CO_rel_f
EB_rel_f
CA_rel_f

HE_evp_f      Reservoir evaporation flow nodes      evp(i)
EV_evp_f
AB_evp_f
CO_evp_f
EB_evp_f
CA_evp_f
/

*****
*      Subsets of all Flow nodes above by class (function)
*****

inflow(i)      Headwater flow nodes      inflow(i)

/      RG-DN_h_f      Rio Grande headwaters at Del Norte gage CO
      Conejos_h_f      Conejos River headwaters-CO
      CBasn_h_f      Closed Basin Project-CO-pumps into RG for RGCompact
      SangDC_h_f      Sangre De Cristo Headwater Flows-NM below Lobatos
      SJCham_h_f      San Juan Chama Interbasin transfer from CO to NM
      Chama_h_f      Rio Chama headwaters near CO-NM state line
      Jemez_h_f      Jemez River headwaters above Alb NM
      Puerco_h_f      Rio Puerco above Socorro NM
      Salado_h_f      Rio Salado below Socorro NM
/

river(i)      River gage measurement nodes      river(i)

/      Lobatos_v_f      Lobatos gauge on RG at CO-NM state line
      Embudo_v_f      Embudo gauge on RG northern NM
      Chamita_v_f      Chamita gauge on Rio Chama northern NM
      Otowi_v_f      Otowi gauge on RG downstream of Chama RG confluence
      Cochit_g_v_f      Cochiti Lake outflow gage below Santa Fe NM
      Acacia_v_f      San Acacia gauge near Socorro NM
      Marcial_v_f      San Marcial gauge below Socorro NM

```

```

EB_g_v_f      Elephant Butte Lake outflow gauge near TorC NM
CA_g_v_f      Caballo Lake outflow gauge near Hatch NM
ElPaso_v_f    Rio Grande flow at El Paso below MX delivery
Quitman_v_f    Rio Grande flow at Fort Quitman TX
/
ag(i)          All agriculture nodes          ag(i)
/
SLV_f         Rio Grande Conservancy ag in SL Valley CO
MRGCD_f       Middle RG Conservancy District ag near Albuq NM
EBID_f        Elephant Butte Irr Dist ag near Las Cruces NM
MXAG_f        Mexican irrigated ag diverted at NM TX MX border
EPAG_f        West TX irrigated Ag near El Paso TX
/
mi(i)         All Muncipal and Industrial (M&I) nodes    mi(i)
/
ABQMI_f       City of Albuquerque NM M&I
EPMI_f        City of El Paso TX M&I
/
divert(i)     Diversion nodes          divert(i)
/
SLV_d_f       ag(i) + mi(i) nodes
ABQMI_d_f
MRGCD_d_f
EBID_d_f
MXAG_d_f
EPMI_d_f
EPAG_d_f
/
adivert(divert)  Diversion nodes per agricultural use    adivert(i)
/
SLV_d_f       Rio Grande Conservancy ag in SL Valley CO
MRGCD_d_f     Middle RG Conservancy District ag near Albuq NM
EBID_d_f      Elephant Butte Irr Dist ag near Las Cruces NM
MXAG_d_f      Mexican ag diversion near El Paso TX
EPAG_d_f      West TX irrigated Ag near El Paso TX
/
mdivert (divert)  Diversion nodes per municipal use    mdivert(i)
/
ABQMI_d_f     Albuquerque
EPMI_d_f      El Paso
/
apply(i)       Application nodes          apply(i)
/
SLV_a_f       same nodes as divert(i)
ABQMI_a_f
MRGCD_a_f
EBID_a_f
MXAG_a_f
EPMI_a_f
EPAG_a_f

```

```

/

aapply(apply)      Application nodes per agricultural use      aapply(i)

/      SLV_a_f      same nodes as divert(i)
      MRGCD_a_f
      EBID_a_f
      MXAG_a_f
      EPAG_a_f
/

mapply (apply)     Application nodes per municipal use      mapply(i)

/      ABQMI_a_f
      EPMI_a_f
/

use(i)             Use (ET consumption) nodes = div nodes      use(i)

/      SLV_u_f      same nodes as divert(i)
      ABQMI_u_f
      MRGCD_u_f
      EBID_u_f
      MXAG_u_f
      EPMI_u_f
      EPAG_u_f
/

ause(use)          Ag use nodes                                ause(i)

/      SLV_u_f      same nodes as divert(i)
      MRGCD_u_f
      EBID_u_f
      MXAG_u_f
      EPAG_u_f
/

muse(use)          M&I use nodes                                muse(i)

/      ABQMI_u_f      same nodes as divert(i)
      EPMI_u_f
/

seep(i)            Seepage nodes                                seep(i)

/      SLV_s_f      same nodes as divert(i)
      ABQMI_s_f
      MRGCD_s_f
      EBID_s_f
      MXAG_s_f
      EPMI_s_f

```

```

    EPAG_s_f
/

aseep(seep)      Seepage per Ag node      aseep(i)

/    SLV_s_f      same nodes as divert(i)
  MRGCD_s_f
  EBID_s_f
  MXAG_s_f
  EPAG_s_f
/

mseep(seep)     Seepage per M&I node     mseep(i)

/    ABQMI_s_f    same nodes as divert(i)
  EPMI_s_f
/

pump(i)         Pumping nodes            pump(i)

/    SLV_p_f      same nodes as divert(i)
  ABQMI_p_f
  MRGCD_p_f
  EBID_p_f
  MXAG_p_f
  EPMI_p_f
  EPAG_p_f
/

netseep(i)      Net seepage nodes        netseep(i)

/    SLV_n_f      same nodes as divert(i)
  ABQMI_n_f
  MRGCD_n_f
  EBID_n_f
  MXAG_n_f
  EPMI_n_f
  EPAG_n_f
/

return(i)       Return (surface) flow nodes  return(i)

/    SLV_r_f      same nodes as divert(i)
  ABQMI_r_f
  MRGCD_r_f
  EBID_r_f
  MXAG_r_f
  EPMI_r_f
  EPAG_r_f
/

areturn(return) Ag return (surface) flow nodes  areturn(i)

```



```

/      SLV_r_f      same nodes as divert(i)
      MRGCD_r_f
      EBID_r_f
      MXAG_r_f
      EPAG_r_f
/

mreturn(return)      M&I return (surface) flow nodes      mreturn(i)

/      ABQMI_r_f      same nodes as divert(i)
      EPMI_r_f
/

gwflow(i)      Groundwater return flow nodes      gwflow(i)

/      SLV_g_f      same nodes as divert(i)
      ABQMI_g_f
      MRGCD_g_f
      EBID_g_f
      MXAG_g_f
      EPMI_g_f
      EPAG_g_f
/

aqflow(i)      Addition to aquifer stock nodes      aqflow(i)

/      SLV_dq_f      same nodes as divert(i)
      ABQMI_dq_f
      MRGCD_dq_f
      EBID_dq_f
      MXAG_dq_f
      EPMI_dq_f
      EPAG_dq_f
/

release(i)      Reservoir to river release flow nodes      release(i)

/      HE_rel_f      Heron (net) reservoir releases to Willow Creek-Rio Chama = outflow - inflow
      EV_rel_f      El Vado reservoir releases to Rio Chama
      AB_rel_f      Abiquiu reservoir releases to Rio Chama
      CO_rel_f      Cochiti reservoir releases to Rio Grande mainstem
      EB_rel_f      Elephant Butte reservoir releases to Rio Grande mainstem
      CA_rel_f      Caballo reservoir releases to Rio Grande mainstem
/

evap(i)      Reservoir evaporation      evap(i)

/      HE_evap_f      Heron reservoir evaporation = fn of annual ave exposed surface area
      EV_evap_f      El Vado reservoir evaporation
      AB_evap_f      Abiquiu reservoir evaporation
      CO_evap_f      Cochiti reservoir evaporation
      EB_evap_f      Elephant Butte reservoir evaporation
      CA_evap_f      Caballo Reservoir evaporation
/

```

```

*****
mu(use)      Municipal and Industrial use nodes
*****

/    ABQMI_u_f      Alb M&I
    EPMI_u_f       El Paso M&I
/

*****
au(use)      Agricultural use nodes
*****

/    SLV_u_f        SL Valley
    MRGCD_u_f      Middle Rio G Cons Distr
    EBID_u_f       EButte Irr Dist
    EPAG_u_f       EPaso Ag
/

*****
ad(divert)   Agricultural divert nodes
*****

/    SLV_d_f        SL Valley
    MRGCD_d_f      Middle Rio G Cons Distr
    EBID_d_f       EButte Irr Dist
    EPAG_d_f       EPaso Ag
/

*****
u           Stocks--location of important nodes on Rio Grande Colorado to Mexico
*****

/    HE_res_s      Reservoir stock nodes                res(u)
    EV_res_s
    AB_res_s
    CO_res_s
    EB_res_s
    CA_res_s

    SLV_aqf_s      Aquifer stock nodes                aqf(u)
    ABQMI_aqf_s
    MRGCD_aqf_s
    EBID_aqf_s
    MXAG_aqf_s
    EPMI_aqf_s
    EPAG_aqf_s
/

*****
*           Stock subsets
*****

res(u)      Reservoir stock nodes                res(u)

```

```

/   HE_res_s   Heron Reservoir (stock) near CO-NM state line
   EV_res_s   El Vado Reservoir downstream of Heron on Chama
   AB_res_s   Abiquiu Reservoir on Chama above Espanola NM
   CO_res_s   Cochiti Reservoir on RGR Mainstem downstream of Santa Fe
   EB_res_s   Elephant Butte Reservoir near T or C NM
   CA_res_s   Caballo Reservoir downstream of T or C NM
/

```

```

aqf(u)           Aquifer stock nodes           aqf(u)

```

```

/   SLV_aqf_s   SLV aquifer supporting irrigated Ag
   ABQMI_aqf_s  Alb aquifer (Lake Superior) supporting M&I pumping
   MRGCD_aqf_s  MRGCD aquifer (near Socorro NM)
   EBID_aqf_s   EBID aquifer (Mesilla Bolson)
   MXAG_aqf_s   Mexico aquifer (Hueco Bolson) supporting Ag
   EPMI_aqf_s   El Paso M&I aquifer (Hueco bolson) supporting M&I pump
   EPAG_aqf_s   El Paso Ag aquifer (Hueco Bolson)
/

```

```

*****

```

```

j  crops Major crops in RGB

```

```

/  Alfalfa
   Barley
   Potato

   Smgrainhay
   Sorghumhay
   Cornsilage

   Pimacotton
   Uplandcotton
   Sprlettuce
   Flettuce
   Fonion
   Msyonion
   Ssonion
   Grainsorghum
   Wheat
   Grchile
   Redchile
   Pecan
/

```

```

k  Irrigation technology

```

```

/ fld           flood irrigation -- heavy water application but low ET
  drp          drip irrigation  -- low water application but high ET
/

```

```

on(j) onion crops only

```

```

/  Fonion
   Msyonion
   Ssonion

```

```
/
ch(j) chile crops only
/ Grchile
  Redchile
/
lt(j) lettuce crops only
/ Sprlettuce
  Flettuce
/
***note: crop grouping due to limitations within data
aajk(aapply,j,k) allowed ag apply crop and technology combinations
/
  SLV_a_f.Alfalfa.fld
  SLV_a_f.Barley.fld
  SLV_a_f.Potato.fld
  MRGCD_a_f.Alfalfa.fld
  MRGCD_a_f.Smgrainhay.fld
  MRGCD_a_f.Sorghumhay.fld
  MRGCD_a_f.Cornsilage.fld
  EBID_a_f.Alfalfa.fld
  EBID_a_f.Pimacotton.fld
  EBID_a_f.Uplandcotton.fld
  EBID_a_f.Sprlettuce.fld
  EBID_a_f.Flettuce.fld
  EBID_a_f.Fonion.fld
  EBID_a_f.Msyonion.fld
  EBID_a_f.Ssonion.fld
  EBID_a_f.Grainsorghum.fld
  EBID_a_f.Wheat.fld
  EBID_a_f.Grchile.fld
  EBID_a_f.Redchile.fld
  EBID_a_f.Pecan.fld
  MXAG_a_f.Alfalfa.fld
  MXAG_a_f.Pimacotton.fld
  MXAG_a_f.Uplandcotton.fld
  MXAG_a_f.Pecan.fld
  EPAG_a_f.Alfalfa.fld
  EPAG_a_f.Pimacotton.fld
  EPAG_a_f.Uplandcotton.fld
  EPAG_a_f.Pecan.fld
  EBID_a_f.Alfalfa.drp
  EBID_a_f.Pimacotton.drp
  EBID_a_f.Uplandcotton.drp
  EBID_a_f.Sprlettuce.drp
```

EBID\_a\_f.Flettuce.drp  
EBID\_a\_f.Fonion.drp  
EBID\_a\_f.Msyonion.drp  
EBID\_a\_f.Ssonion.drp  
EBID\_a\_f.Grainsorghum.drp  
EBID\_a\_f.Wheat.drp  
EBID\_a\_f.Grchile.drp  
EBID\_a\_f.Redchile.drp  
EBID\_a\_f.Pecan.drp

/

aujk(ause,j,k) *Allowed ag use crop and technology combinations*

/

SLV\_u\_f.Alfalfa.fld  
SLV\_u\_f.Barley.fld  
SLV\_u\_f.Potato.fld

MRGCD\_u\_f.Alfalfa.fld  
MRGCD\_u\_f.Smgrainhay.fld  
MRGCD\_u\_f.Sorghumhay.fld  
MRGCD\_u\_f.Cornsilage.fld

EBID\_u\_f.Alfalfa.fld  
EBID\_u\_f.Pimacotton.fld  
EBID\_u\_f.Uplandcotton.fld  
EBID\_u\_f.Sprlettuce.fld  
EBID\_u\_f.Flettuce.fld  
EBID\_u\_f.Fonion.fld  
EBID\_u\_f.Msyonion.fld  
EBID\_u\_f.Ssonion.fld  
EBID\_u\_f.Grainsorghum.fld  
EBID\_u\_f.Wheat.fld  
EBID\_u\_f.Grchile.fld  
EBID\_u\_f.Redchile.fld  
EBID\_u\_f.Pecan.fld

MXAG\_u\_f.Alfalfa.fld  
MXAG\_u\_f.Pimacotton.fld  
MXAG\_u\_f.Uplandcotton.fld  
MXAG\_u\_f.Pecan.fld

EPAG\_u\_f.Alfalfa.fld  
EPAG\_u\_f.Pimacotton.fld  
EPAG\_u\_f.Uplandcotton.fld  
EPAG\_u\_f.Pecan.fld

EBID\_u\_f.Alfalfa.drp  
EBID\_u\_f.Pimacotton.drp  
EBID\_u\_f.Uplandcotton.drp  
EBID\_u\_f.Sprlettuce.drp  
EBID\_u\_f.Flettuce.drp  
EBID\_u\_f.Fonion.drp  
EBID\_u\_f.Msyonion.drp  
EBID\_u\_f.Ssonion.drp  
EBID\_u\_f.Grainsorghum.drp

EBID\_u\_f.Wheat.drp  
EBID\_u\_f.Grchile.drp  
EBID\_u\_f.Redchile.drp  
EBID\_u\_f.Pecan.drp

/

asjk(aseep,j,k) *Allowed ag seep crop and technology combinations*

/

SLV\_s\_f.Alfalfa.fld  
SLV\_s\_f.Barley.fld  
SLV\_s\_f.Potato.fld

MRGCD\_s\_f.Alfalfa.fld  
MRGCD\_s\_f.Smgrainhay.fld  
MRGCD\_s\_f.Sorghumhay.fld  
MRGCD\_s\_f.Cornsilage.fld

EBID\_s\_f.Alfalfa.fld  
EBID\_s\_f.Pimacotton.fld  
EBID\_s\_f.Uplandcotton.fld  
EBID\_s\_f.Sprlettuce.fld  
EBID\_s\_f.Flettuce.fld  
EBID\_s\_f.Fonion.fld  
EBID\_s\_f.Msyonion.fld  
EBID\_s\_f.Ssonion.fld  
EBID\_s\_f.Grainsorghum.fld  
EBID\_s\_f.Wheat.fld  
EBID\_s\_f.Grchile.fld  
EBID\_s\_f.Redchile.fld  
EBID\_s\_f.Pecan.fld

MXAG\_s\_f.Alfalfa.fld  
MXAG\_s\_f.Pimacotton.fld  
MXAG\_s\_f.Uplandcotton.fld  
MXAG\_s\_f.Pecan.fld

EPAG\_s\_f.Alfalfa.fld  
EPAG\_s\_f.Pimacotton.fld  
EPAG\_s\_f.Uplandcotton.fld  
EPAG\_s\_f.Pecan.fld

EBID\_s\_f.Alfalfa.drp  
EBID\_s\_f.Pimacotton.drp  
EBID\_s\_f.Uplandcotton.drp  
EBID\_s\_f.Sprlettuce.drp  
EBID\_s\_f.Flettuce.drp  
EBID\_s\_f.Fonion.drp  
EBID\_s\_f.Msyonion.drp  
EBID\_s\_f.Ssonion.drp  
EBID\_s\_f.Grainsorghum.drp  
EBID\_s\_f.Wheat.drp  
EBID\_s\_f.Grchile.drp  
EBID\_s\_f.Redchile.drp  
EBID\_s\_f.Pecan.drp

/

arjk(areturn,j,k) *Allowed ag return crop and technology combinations*

/

SLV\_r\_f.Alfalfa.fld  
SLV\_r\_f.Barley.fld  
SLV\_r\_f.Potato.fld

MRGCD\_r\_f.Alfalfa.fld  
MRGCD\_r\_f.Smgrainhay.fld  
MRGCD\_r\_f.Sorghumhay.fld  
MRGCD\_r\_f.Cornsilage.fld

EBID\_r\_f.Alfalfa.fld  
EBID\_r\_f.Pimacotton.fld  
EBID\_r\_f.Uplandcotton.fld  
EBID\_r\_f.Sprlettuce.fld  
EBID\_r\_f.Flettuce.fld  
EBID\_r\_f.Fonion.fld  
EBID\_r\_f.Msyonion.fld  
EBID\_r\_f.Ssonion.fld  
EBID\_r\_f.Grainsorghum.fld  
EBID\_r\_f.Wheat.fld  
EBID\_r\_f.Grchile.fld  
EBID\_r\_f.Redchile.fld  
EBID\_r\_f.Pecan.fld

MXAG\_r\_f.Alfalfa.fld  
MXAG\_r\_f.Pimacotton.fld  
MXAG\_r\_f.Uplandcotton.fld  
MXAG\_r\_f.Pecan.fld

EPAG\_r\_f.Alfalfa.fld  
EPAG\_r\_f.Pimacotton.fld  
EPAG\_r\_f.Uplandcotton.fld  
EPAG\_r\_f.Pecan.fld

EBID\_r\_f.Alfalfa.drp  
EBID\_r\_f.Pimacotton.drp  
EBID\_r\_f.Uplandcotton.drp  
EBID\_r\_f.Sprlettuce.drp  
EBID\_r\_f.Flettuce.drp  
EBID\_r\_f.Fonion.drp  
EBID\_r\_f.Msyonion.drp  
EBID\_r\_f.Ssonion.drp  
EBID\_r\_f.Grainsorghum.drp  
EBID\_r\_f.Wheat.drp  
EBID\_r\_f.Grchile.drp  
EBID\_r\_f.Redchile.drp  
EBID\_r\_f.Pecan.drp

/

\*\*\*\*\*  
t *Time*  
\*\*\*\*\*

```

/ 2006*2010      years 2006 - 2010  /

tlast(t)  Terminal period among all periods above
;

tlast(t) = yes $(ord(t) eq card(t));  > GAMS language -- picks last pd
;

* -----
* Some mathematical tricks below streamline the model
* -----

* lets some tables' nodes be rows or columns

ALIAS (i,ip);
ALIAS (t,tp);
ALIAS (river, riverp);
ALIAS (divert, divertp);

* tables below defined as identity matrices, to compare different but equivalent nodes
* e.g. a divert node is the same place as an apply, use, return, pump node

PARAMETER
ID_au(apply, use)  identity matrix connects apply nodes to use nodes
ID_pu(pump, use)  identity matrix connects pump nodes to use nodes
ID_pa(pump, apply) identify matrix connects pump nodes to apply nodes
ID_ap(apply, pump) identity matrix connects apply nodes to pump nodes
ID_ad(apply,divert) identity matrix connects apply nodes to divert nodes
ID_ua(ause, aapply) identity matrix connects ag use nodes to ag apply nodes
ID_us(ause, aseep) identity matrix connects ag use nodes to ag seep nodes
ID_ur(ause,areturn) identity matrix connects ag use nodes to ag return nodes
;

ID_au(apply, use) $ (ord(apply) eq ord(use )) = 1;
ID_pu(pump, use) $ (ord(pump ) eq ord(use )) = 1;
ID_pa(pump, apply) $ (ord(pump ) eq ord(apply )) = 1;
ID_ap(apply, pump) $ (ord(apply) eq ord(pump )) = 1;
ID_ad(apply,divert) $ (ord(apply) eq ord(divert )) = 1;

ID_ua(ause, aapply) $ (ord(ause ) eq ord(aapply )) = 1;
ID_us(ause, aseep) $ (ord(ause ) eq ord(aseep )) = 1;
ID_ur(ause,areturn) $ (ord(ause ) eq ord(areturn)) = 1;

*-----

***** Section 2 *****
* This section defines all data in 3 formats *
* 1. Scalars (single numbers), *
* 2. Parameters (columns of numbers) or *
* 3. Tables (data in rows and columns) *
*****

* Below are several maps summarizing a basin's geometry
* By geometry we mean location of mainstems, tributaries, confluence,
* source nodes, use nodes, return flow nodes, reservoir nodes, etc.

```



\* Basin geometry is summarized through judicious use of numbers 1, -1, and 0 (blank)

\*\*\*\*\*

\* Map #1:

\* Each column below is a streamgage. Each row is a source or use of water.  
 \* Flow at ea gage (column) is directly influenced by at least 1 upstream row.  
 \* SOURCE adds to columns flow (+1)  
 \* USE deplete from col flow (-1)  
 \* BLANK has no effect on col flow ( )  
 \* Geometry accounts for all sources (supplies) and uses (demands) in basin

\* Map is used to produce coefficients in equations below to define:  
 $X(\text{river}) = B_{hv} * X(\text{inflow}) + B_{vv} * X(\text{river}) + B_{dv} * X(\text{divert})$   
 $+ B_{rv} * X(\text{return}) + B_{gv} * X(\text{gwflows}) + B_{Lr} * X(\text{rel})$

\* These B coeff vectors are stacked below as a single matrix, Bv

\*\*\*\*\*

TABLE Bv(i,river) Hydrologic Balance Table

\*\*\*\*\* Column Heads are River Gauges \*\*\*\*\*

Lobatos\_v\_f Embudo\_v\_f Chamita\_v\_f Otowi\_v\_f Coch\_g\_v\_f Acacia\_v\_f Marcial\_v\_f EB\_g\_v\_f CA\_g\_v\_f ElPaso\_v\_f Quitman\_v\_f

\* ----- headwater inflow node rows (+) -----

RG-DN_h_f	1									
Conejos_h_f	1									
CBasn_h_f	1									
SangDC_h_f		1								
SJCham_h_f			1							
Chama_h_f			1							
Jemez_h_f					1					
Puerco_h_f					1					
Salado_h_f							1			

\* ----- river gage node rows (+) -----

Lobatos_v_f	1									
Embudo_v_f			1							
Chamita_v_f			1							
Otowi_v_f				1						
Coch_g_v_f					1					
Acacia_v_f						1				
Marcial_v_f							1			
EB_g_v_f								1		
CA_g_v_f									1	
ElPaso_v_f										1
Quitman_v_f										

\* ----- diversion nodes (-) -----

SLV_d_f	-1									
ABQMI_d_f					-1					
MRGCD_d_f					-1					
EBID_d_f									-1	
MXAG_d_f									-1	
EPMI_d_f									-1	
EPAG_d_f										-1

\* ----- return flow node rows (+) -----

```

SLV_r_f          1
ABQMI_r_f
MRGCD_r_f
EBID_r_f
MXAG_r_f
EPMI_r_f
EPAG_r_f
* ----- groundwater inflow node rows (+) -----
SLV_g_f          1
ABQMI_g_f
MRGCD_g_f
EBID_g_f
MXAG_g_f
EPMI_g_f
EPAG_g_f
* ----- reservoir release (outflow) to river -- stock-to-flow rows (+) -----
HE_rel_f          1
EV_rel_f          1
AB_rel_f          1
CO_rel_f          1
EB_rel_f          1
CA_rel_f          1
;
*****
* Map #2:
* Enforces nonnegative flows at each use node (wet river)
* water sources are rows. Diversion nodes are columns.
* For any column, diversion < summed flows from upstream sources (rows)
* e.g. SLV Colorado ag use < flows from RG and Conejos headwater sources
*
*  $X(\text{divert}) < B_{hd} * X(\text{inflow}) + B_{rd} * X(\text{river}) + B_{dd} * X(\text{divert}) +$ 
*  $B_{rd} * X(\text{return}) + B_{gd} * X(\text{gwflow}) + B_{ld} * X(\text{rel})$ 
*
* These B coeff vectors are stacked below as the matrix, Bd
*****
TABLE Bd(i, divert)  Wet river table
* ----- Col Heads are Diversion nodes -----
          SLV_d_f      ABQMI_d_f      MRGCD_d_f      EBID_d_f      MXAG_d_f      EPMI_d_f      EPAG_d_f
* ----- headwater inflow nodes (+)-----
RG-DN_h_f          1
Conejos_h_f        1
CBasn_h_f
SangDC_h_f
SJCham_h_f
Chama_h_f
Jemez_h_f          1
Puerco_h_f         1
Salado_h_f
* ----- river gage nodes -----
Lobatos_v_f
Embudo_v_f

```

```

Chamita_v_f
Otowi_v_f
Coch_g_v_f          1      1
Acacia_v_f
Marcial_v_f
EB_g_v_f
CA_g_v_f            1      1      1      1
ElPaso_v_f
Quitman_v_f
* ----- diversion nodes (-) -----
SLV_d_f
ABQMI_d_f           -1
MRGCD_d_f
EBID_d_f            -1      -1
MXAG_d_f            -1
EPMI_d_f
EPAG_d_f
* ----- return flow nodes (+) -----
SLV_r_f
ABQMI_r_f           1
MRGCD_r_f
EBID_r_f            1      1
MXAG_r_f            1
EPMI_r_f
EPAG_r_f
* ----- groundwater inflow nodes (+) -----
SLV_g_f
ABQMI_g_f           1
MRGCD_g_f
EBID_g_f            1      1
MXAG_g_f            1
EPMI_g_f
EPAG_g_f
* ----- reservoir outflow stock-to-flow node row (+)-----
HE_rel_f
EV_rel_f
AB_rel_f
CO_rel_f
EB_rel_f
CA_rel_f
* -----

```

\* Map #3

\* Defines pan evaporation in inches - translate to af lost per per exposed acre per year by reservoir

TABLE Be(evap, res)

	HE_res_s	EV_res_s	AB_res_s	CO_res_s	EB_res_s	CA_res_s
HE_evp_f	56.0					
EV_evp_f		56.0				
AB_evp_f			56.0			
CO_evp_f				93.0		
EB_evp_f					111.2	
CA_evp_f						111.2

;

Be(evap,res) = Be(evap,res)/12; > inches to feet

\*display Be;

\* -----

\*\*\*\*\*

\*\*\*\*\*

\* Map #4:

\* Defines water applied as diversion plus pumping

\*  $X(\text{apply}) = B_{da} * X(\text{divert}) + B_{pa} * X(\text{pump})$

\* These two B coefficient vectors are stacked below as the matrix, Ba

TABLE Ba(i, apply) Table defines water applied

\* ----- Apply nodes -----

SLV\_a\_f ABQMI\_a\_f MRGCD\_a\_f EBID\_a\_f MXAG\_a\_f EPMI\_a\_f EPAG\_a\_f

\* ----- divert nodes (+) -----

SLV_d_f	1						
ABQMI_d_f		1					
MRGCD_d_f			1				
EBID_d_f				1			
MXAG_d_f					1		
EPMI_d_f						1	
EPAG_d_f							1

\* ----- pumping nodes (+) -----

SLV_p_f	1						
ABQMI_p_f		1					
MRGCD_p_f			1				
EBID_p_f				1			
MXAG_p_f					1		
EPMI_p_f						1	
EPAG_p_f							1

\*-----

;

\*\*\*\*\*

\* Map #5:

\* Defines use as a constant proportion, Bau, of water applied

\*  $X(\text{use}) = B_{au} * X(\text{apply})$

\* This coeff vector is the matrix, Bu

\*\*\*\*\*

TABLE Bu(apply, use) Table defines consumptive use

\* Best data for next 3 tables may be in Papodolous report (1996?). Generally, these data  
\* are scarce, scattered, and hard to validate. Many questions in basic science and most  
\* questions in applied science regarding groundwater pumping connections to surface supply  
\* remain unanswered. For relevant theory, the best article may be a December 1940 paper by

\* M King-Hubbert, The Theory of Groundwater Motion, Journal of Geology, pp 785-944 (160 pp)

\* MX data are educated guesses

\* Albuquerque return flows ABQ Water home page -- <http://www.cabq.gov/waterresources/sjc.html>

```
* ----- Use nodes -----
*          SLV_u_f   ABQMI_u_f   MRGCD_u_f   EBID_u_f   MXAG_u_f   EPMI_u_f   EPAG_u_f
* ----- apply nodes (+) -----
SLV_a_f      0.50
ABQMI_a_f           0.441
MRGCD_a_f                0.25
EBID_a_f                    0.46
MXAG_a_f                        0.46
EPMI_a_f                            0.441
EPAG_a_f                                0.46
```

PARAMETER Bu\_p(use) *Translates Bu above into a parameter with one value at each use node*

Bu\_p(use) = sum(apply, Bu(apply,use));

\*\*\*\*\*

\* Map #6:

\* Defines seepage as a constant proportion, Bas, of water applied

\*  $X(\text{seep}) = Bas * X(\text{apply})$

\* This coeff vector is the matrix, Bs

\*\*\*\*\*

TABLE Bs(apply, seep) *Table defines seepage to groundwater aquifer*

```
* ----- Seepage nodes -----
*          SLV_s_f   ABQMI_s_f   MRGCD_s_f   EBID_s_f   MXAG_s_f   EPMI_s_f   EPAG_s_f
* ----- apply nodes (+) -----
SLV_a_f      0.50
ABQMI_a_f           0.059
MRGCD_a_f                0.19
EBID_a_f                    0.34
MXAG_a_f                        0.34
EPMI_a_f                            0.059
EPAG_a_f                                0.34
```

;

\*\*\*\*\*

\* Map #7:

\* Defines return flow as a constant proportion, Bar, of water applied

\*  $X(\text{return}) = Bar * X(\text{apply})$

\* This coeff vector is the matrix, Br

\*\*\*\*\*

TABLE Br(apply, return) *Table defines return flow to river (surface flow)*

```

* ----- Return flow nodes -----
*          SLV_r_f  ABQMI_r_f  MRGCD_r_f  EBID_r_f  MXAG_r_f  EPMI_r_f  EPAG_r_f
* ----- apply nodes (+) -----
SLV_a_f      0.00
ABQMI_a_f           0.500
MRGCD_a_f           0.56
EBID_a_f           0.20
MXAG_a_f           0.20
EPMI_a_f           0.500
EPAG_a_f           0.20
*-----
;

```

PARAMETER APPLY\_FLOW\_ALLOC(apply) *check: flow allocations should sum to 1*

```

;
APPLY_FLOW_ALLOC(apply) =  sum(use,   Bu(apply, use  ))
                          +  sum(seep,  Bs(apply, seep ))
                          +  sum(return, Br(apply, return));

```

*\*display apply\_flow\_alloc;*

\*\*\*\*\*

*\* Map #8:*

*\* Table defines net seepage as seepage minus pumping*

*\**

*\* Tabled entries = proportion seepage and pumping flow by net seepage column nodes*

*\* (+) means the row adds to the column's net seepage*

*\* (-) means the row subtracts from column's net seepage*

*\* ( ) no effect*

*\*  $X(\text{netseep}) = B_{sn} * X(\text{seep}) + B_{pn} * X(\text{pump})$*

\*\*\*\*\*

*\* These two B coefficient vectors are stacked below as the matrix, Bn*

TABLE Bn(i, netseep) *Table defines net seepage to (+) or from (-) aquifer*

\*\*\*\*\* Column Heads are Net Seepage Nodes \*\*\*\*\*

```

          SLV_n_f  ABQMI_n_f  MRGCD_n_f  EBID_n_f  MXAG_n_f  EPMI_n_f  EPAG_n_f
* ----- seepage node rows (+) -----
SLV_s_f      1
ABQMI_s_f           1
MRGCD_s_f           1
EBID_s_f           1
MXAG_s_f           1
EPMI_s_f           1
EPAG_s_f           1
* ----- pump node rows (-) -----
SLV_p_f      -1
ABQMI_p_f           -1
MRGCD_p_f           -1
EBID_p_f           -1

```

```

MXAG_p_f          -1
EPMI_p_f          -1
EPAG_p_f          -1
-----

```

```

;

```

```

*****

```

```

* Best data for next 2 tables may be in Papodolous report (1996?). They allocate
* groundwater flow from net seepage to 2 sources-sinks: (1) river and (2) aquifer.
* Net seepage hydrologically farther from river has a smaller pct affecting river.

```

```

* Like the tables above that allocate water application among use, seepage, and return
* flows, seepage allocation data are scarce, scattered, and hard to validate. Many
* questions in basic science and most in applied science regarding groundwater pumping
* connections to surface supply v. aquifer drawdowns remain unanswered. For relevant theory,
* the best article may be a Dec 1940 paper by M King-Hubbert, The Theory of Groundwater
* Motion, J. of Geology, pp 785-944 (160 pages, yes 160 pages total)

```

```

* Adding downward, coefficients from next two tables should also add to 1.0, i.e. for each
* diversion node, total groundwater flow percentage summed from the river and from the
* aquifer should add to 1.0;

```

```

*****

```

```

* Map #9:

```

```

* Table defines groundwater flow to river as a fn of net seepage
*
* Tabled entries = proportion net seepage contributing to groundwater flow column nodes
* (+) means the row adds to the column's gw flow
* (-) means the row subtracts from column's gw flow
* ( ) no effect

```

```

* X(gwflow) = Bng * X(netseep)
*****

```

```

TABLE Bg(netseep, gwflow) Table defines groundwater flow to (+) or from (-) river

```

```

***** Column Heads are Groundwater Flow Nodes *****

```

	SLV_g_f	ABQMI_g_f	MRGCD_g_f	EBID_g_f	MXAG_g_f	EPMI_g_f	EPAG_g_f
* ----- net seepage node rows (+) -----							
SLV_n_f	0						
ABQMI_n_f	0.60						
MRGCD_n_f		0.60					
EBID_n_f			0.40				
MXAG_n_f				0.80			
EPMI_n_f					0.80		
EPAG_n_f						0.80	

```

;

```

```

*****

```

```

* Map #10: complements Map 8. Sum of Bg and Bq coeffs for any colum = 1 for mass balance

```

```

* Table defines added aquifer volume as a fn of net seepage
*

```

```

* Tabled entries = proportion net seepage contributing to added aquifer volume
* (+) means the row adds to the column's aquifer volume (change in aq stock = flow)
* (-) means the row subtracts from column's aquifer volume
* ( ) no effect

```

```

* X(aqflow) = Bqg * X(netseep)

```

```

*****

```

```

TABLE Bq(netseep, aqflow) Table defines gain (+) or loss (-) in aquifer volume

```

```

***** Column Heads are Aquifer volume change (flow) nodes *****

```

	SLV_dq_f	ABQMI_dq_f	MRGCD_dq_f	EBID_dq_f	MXAG_dq_f	EPMI_dq_f	EPAG_dq_f
SLV_n_f	1						
ABQMI_n_f		0.40					
MRGCD_n_f			0.40				
EBID_n_f				0.60			
MXAG_n_f					0.20		
EPMI_n_f						0.20	
EPAG_n_f							0.20

```

PARAMETER SEEP_FLOW_ALLOC(netseep) check: allocations of net seepages should sum to 1

```

```

;
SEEP_FLOW_ALLOC(netseep) = sum(gwflow, Bg(netseep, gwflow))
+ sum(aqflow, Bq(netseep, aqflow));

```

```

*DISPLAY SEEP_FLOW_ALLOC;

```

```

*****

```

```

* Next two maps link stocks (res and aquifers) to flows (release and net seepage)
*****
* Map #11:

```

```

* Table relates reserv stocks in a pd to its prev pd' stocks minus net releases.
* For any reservoir stock node at the column head
* (-1) :added water at flow node -- thru releases -- subtracts from column's res stock
* ( ) :added water at flow node has no effect on column's reservoir stock

```

```

* Z(res(t)) = Z(res(t-1)) + BLr * X(release(t))

```

```

*****

```

```

TABLE BLr(release, res) Links reservoir releases to downstream flows

```

```

***** Column Heads are Reservoir Stocks -- rows are release flows *****
***** Table = diagonal matrix for > 1 reservoir -- only 1 for now *****

```

	HE_res_s	EV_res_s	AB_res_s	CO_res_s	EB_res_s	CA_res_s
HE_rel_f	-1					
EV_rel_f		-1				
AB_rel_f			-1			
CO_rel_f				-1		
EB_rel_f					-1	
CA_rel_f						-1

```

-----

```



\*\*\*\*\*

\* Map #12:

\* Table relates reservoir stocks to evaporation  
\* (-1): added evap subtracts a reservoir's volume  
\* ( ): added evap has no effect on a reservoir's vol

TABLE Ber(evap, res) Links reservoir evaporation to volume loss

\*\*\*\*\* Column Heads are reservoir stocks -- rows are evaporation loss flows \*\*\*\*\*  
\*\*\*\*\* Table = diagonal matrix for > 1 reservoir -- only 1 for now \*\*\*\*\*

	HE_res_s	EV_res_s	AB_res_s	CO_res_s	EB_res_s	CA_res_s
HE_evap_f	-1					
EV_evap_f		-1				
AB_evap_f			-1			
CO_evap_f				-1		
EB_evap_f					-1	
CA_evap_f						-1

-----

\*\*\*\*\*  
\* Map #13:

\* Table relates aquifer volume stock in a period to its prev periods' stocks plus  
\* net seepage (seepage minus pumping)  
\* For any aquifer stock node at the column head  
\* (+1):added water at flow (net seep) node -- thru net seepage -- adds to col aqf stck  
\* ( ):added water at pump has no effect on column's aqf stock

$$Z(aqf(t)) = Z(aqf(t-1)) + BQv * X(aqflow(t))$$

TABLE BQv(aqflow, aqf) Links aquifer pumping to aquifer volume and depth

\*\*\*\*\* Column Heads are Aquifer stocks -- rows are pumping flows \*\*\*\*\*  
\*\*\*\*\* Table = diagonal matrix for > 1 aquifers \*\*\*\*\*

	SLV_aqf_s	ABQMI_aqf_s	MRGCD_aqf_s	EBID_aqf_s	MXAG_aqf_s	EPMI_aqf_s	EPAG_aqf_s
SLV_dq_f	1						
ABQMI_dq_f		1					
MRGCD_dq_f			1				
EBID_dq_f				1			
MXAG_dq_f					1		
EPMI_dq_f						1	
EPAG_dq_f							1

\*\*\*\*\*  
\* END OF BASIN HYDROLOGY MAPS \*  
\*\*\*\*\*

\*\*\*\*\*  
\* AGRICULTURAL (CROP) MAPS BEGIN \*  
\*\*\*\*\*

\*\*\*\*\*

\* Crop Map #1: CROP WATER USE BY CROP AND IRRIGATION TECHNOLOGY

\*\*\*\*\*

\* Table defines hydrologic outcomes after applying water to crops

\*

\* Tabled entries = quantity of water allocated from total water applied

\* entry = the row required water per acre of column's crop produced

\* Top matrix below is water per acre applied by crops at ag nodes

\* Three B coefficient vectors are stacked beneath top matrix.

\* Top matrix (water applied) allocates water to 3 bottom matrices

\* Water apply = water use + water seep + water return.

\* The whole matrix is defined as BBa\_p

TABLE BBa\_p(i, j, k) Per acre crop water applied for various use indicators

\*\*\*\*\* Column Heads are Crops \*\*\*\*\*

	Alfalfa.fld	Barley.fld	Potato.fld	Smgrainhay.fld	Sorghumhay.fld	Cornsilage.fld			
*----- apply node rows (+) -----*									
SLV_a_f	2.9	3.33	3.33	2.0	2.0	3.5			
MRGCD_a_f	3.67	3.33	3.33	2.0	2.0	3.5			
EBID_a_f	5.0	3.33	3.33	2.0	2.0	3.5			
MXAG_a_f	5.0	3.33	3.33	2.0	2.0	3.5			
EPAG_a_f	5.0	3.33	3.33	2.0	2.0	3.5			
*----- use node rows (+) -----*									
SLV_u_f	1.25	1.43	1.43	0.86	0.86	1.5			
MRGCD_u_f	1.58	1.43	1.43	0.86	0.86	1.5			
EBID_u_f	2.15	1.43	1.43	0.86	0.86	1.5			
MXAG_u_f	2.15	1.43	1.43	0.86	0.86	1.5			
EPAG_u_f	2.15	1.43	1.43	0.86	0.86	1.5			
*----- seep node rows (+) -----*									
SLV_s_f	1.65	1.9	1.9	1.14	1.14	2.0			
MRGCD_s_f	1.045	0.95	0.95	0.57	0.57	1.0			
EBID_s_f	1.425	0.95	0.95	0.57	0.57	1.0			
MXAG_s_f	1.425	0.95	0.95	0.57	0.57	1.0			
EPAG_s_f	1.425	0.95	0.95	0.57	0.57	1.0			
*----- return flow node rows (+) -----*									
SLV_r_f	0.0	0.0	0.0	0.0	0.0	0.0			
MRGCD_r_f	1.045	0.95	0.95	0.57	0.57	1.0			
EBID_r_f	1.425	0.95	0.95	0.57	0.57	1.0			
MXAG_r_f	1.425	0.95	0.95	0.57	0.57	1.0			
EPAG_r_f	1.425	0.95	0.95	0.57	0.57	1.0			
*-----*									
+ Pimacotton.fld Uplandcotton.fld Sprlettuce.fld Flettuce.fld Fonion.fld Msyonion.fld Ssonion.fld									
*----- apply node rows (+) -----*									
SLV_a_f	2.75	2.75	2.5	3.33	4.67	4.0	4.75		
MRGCD_a_f	2.75	2.75	2.5	3.33	4.67	4.0	4.75		
EBID_a_f	2.75	2.75	2.5	3.33	4.67	4.0	4.75		
MXAG_a_f	2.75	2.75	2.5	3.33	4.67	4.0	4.75		
EPAG_a_f	2.75	2.75	2.5	3.33	4.67	4.0	4.75		
*----- use node rows (+) -----*									
SLV_u_f	1.18	1.18	1.08	1.43	2.0	2.28	2.71		
MRGCD_u_f	1.18	1.18	1.08	1.43	2.0	2.28	2.71		
EBID_u_f	1.18	1.18	1.08	1.43	2.0	2.28	2.71		

MXAG_u_f	1.18	1.18	1.08	1.43	2.0	2.28	2.71
EPAG_u_f	1.18	1.18	1.08	1.43	2.0	2.28	2.71
*----- seep node rows (+) -----*							
SLV_s_f	1.57	1.57	1.42	1.9	2.67	1.72	2.04
MRGCD_s_f	0.785	0.785	0.71	0.95	1.335	0.86	1.02
EBID_s_f	0.785	0.785	0.71	0.95	1.335	0.86	1.02
MXAG_s_f	0.785	0.785	0.71	0.95	1.335	0.86	1.02
EPAG_s_f	0.785	0.785	0.71	0.95	1.335	0.86	1.02
*----- return flow node rows (+) -----*							
SLV_r_f	0.0	0.0	0.0	0.0	0.0	0.0	0.0
MRGCD_r_f	0.785	0.785	0.71	0.95	1.335	0.86	1.02
EBID_r_f	0.785	0.785	0.71	0.95	1.335	0.86	1.02
MXAG_r_f	0.785	0.785	0.71	0.95	1.335	0.86	1.02
EPAG_r_f	0.785	0.785	0.71	0.95	1.335	0.86	1.02

+	Grainsorghum.fld	Wheat.fld	Grchile.fld	Redchile.fld	Pecan.fld
*----- apply node rows (+) -----*					
SLV_a_f	2.0	2.5	4.6	5.0	6.0
MRGCD_a_f	2.0	2.5	4.6	5.0	6.0
EBID_a_f	2.0	2.5	4.6	5.0	6.0
MXAG_a_f	2.0	2.5	4.6	5.0	6.0
EPAG_a_f	2.0	2.5	4.6	5.0	6.0
*----- use node rows (+) -----*					
SLV_u_f	0.86	1.075	1.98	2.15	2.58
MRGCD_u_f	0.86	1.075	1.98	2.15	2.58
EBID_u_f	0.86	1.075	1.98	2.15	2.58
MXAG_u_f	0.86	1.075	1.98	2.15	2.58
EPAG_u_f	0.86	1.075	1.98	2.15	2.58
*----- seep node rows (+) -----*					
SLV_s_f	1.14	1.425	2.62	2.85	3.42
MRGCD_s_f	0.57	0.7125	1.31	1.425	1.71
EBID_s_f	0.57	0.7125	1.31	1.425	1.71
MXAG_s_f	0.57	0.7125	1.31	1.425	1.71
EPAG_s_f	0.57	0.7125	1.31	1.425	1.71
*----- return flow node rows (+) -----*					
SLV_r_f	0.0	0.0	0.0	0.0	0.0
MRGCD_r_f	0.57	0.7125	1.31	1.425	1.71
EBID_r_f	0.57	0.7125	1.31	1.425	1.71
MXAG_r_f	0.57	0.7125	1.31	1.425	1.71
EPAG_r_f	0.57	0.7125	1.31	1.425	1.71

\*\*note: starts table for EBID drip irrigation

+	Alfalfa.drp	Pimacotton.drp	Uplandcotton.drp	Sprlettuce.drp	Flettuce.drp	Fonion.drp	Msyonion.drp	Ssonion.drp
*----- apply node rows (+) -----*								
EBID_a_f	1.43	0.65	0.65	0.86	0.78	1.5	1.5	1.5
*----- use node rows (+) -----*								
EBID_u_f	1.43	0.65	0.65	0.86	0.78	1.5	1.5	1.5
*----- seep node rows (+) -----*								
EBID_s_f	0.00	0.00	0.00	0.00	0.00	0.0	0.0	0.0
*----- return flow node rows (+) -----*								
EBID_r_f	0.00	0.00	0.00	0.00	0.00	0.0	0.0	0.0

```

+           Grainsorghum.drp  Wheat.drp  Grchile.drp  Redchile.drp  Pecan.drp
*----- apply node rows (+) -----
EBID_a_f      0.65      0.65      1.5          1.5          1.72
*----- use node rows (+) -----
EBID_u_f      0.65      0.65      1.5          1.5          1.72
*----- seep node rows (+) -----
EBID_s_f      0.0        0.0        0.0          0.0          0.00
*----- return flow node rows (+) -----
EBID_r_f      0.0        0.0        0.0          0.0          0.00
*-----
;

```

\* Distribution of applied water (for seep and return data) obtained with a phone conversation with David Gensler of the  
 \* MRGCD general office in Alb, NM.  
 \* Distribution in MRGCD is as follows: 100% apply, 43% ET, 27.5% seep, 27.5% return flow.  
 \* Based on surface water infrastructure flow gauges.  
 \* This distribution is currently being applied to all nodes until more node-specific ET data becomes available.  
 \* MRGCD data applied to all Ag nodes except SLV which returns no flows to river, all seepage.  
 \* SLV: 43% crop ET, 57% seepage, 0% return to river.

\*DISPLAY BBA\_p;

\*\*\*\*\*  
 \* AGRICULTURE MAP #2 CROP YIELDS BY CROP AND IRRIGATION TECHNOLOGY  
 \*\*\*\*\*

TABLE Yield\_p(use, j, k) Expected yields

\*\*\*\*\* Column Heads are Crops \*\*\*\*\*

	Alfalfa.fld	Barley.fld	Potato.fld	Smgrainhay.fld	Sorghumhay.fld	Cornsilage.fld			
SLV_u_f	5.0	3.26	310.0	0.0	0.0	0.0			
MRGCD_u_f	5.0	0.0	0.0	2.5	2.0	20.0			
EBID_u_f	8.0	0.0	0.0	0.0	0.0	0.0			
MXAG_u_f	8.0	0.0	0.0	0.0	0.0	0.0			
EPAG_u_f	8.0	0.0	0.0	0.0	0.0	0.0			

  

	Pimacotton.fld	Uplandcotton.fld	Sprlettuce.fld	Flettuce.fld	Fonion.fld	Msyonion.fld	Ssonion.fld
SLV_u_f	0.0	0.0	0.0	0.0	0.0	0.0	0.0
MRGCD_u_f	0.0	0.0	0.0	0.0	0.0	0.0	0.0
EBID_u_f	750.0	1000.0	475.0	500.0	1200.0	675.0	825.0
MXAG_u_f	750.0	1000.0	0.0	0.0	0.0	0.0	0.0
EPAG_u_f	750.5	1000.0	0.0	0.0	0.0	0.0	0.0

  

	Grainsorghum.fld	Wheat.fld	Grchile.fld	Redchile.fld	Pecan.fld
SLV_u_f	0.0	0.0	0.0	0.0	0.0
MRGCD_u_f	0.0	0.0	0.0	0.0	0.0
EBID_u_f	40.0	92.0	11.0	3500.0	1158.14
MXAG_u_f	0.0	0.0	0.0	0.0	1158.14
EPAG_u_f	0.0	0.0	0.0	0.0	1158.14

\*\*note: starts EBID DI yields

	Alfalfa.drp	Pimacotton.drp	Uplandcotton.drp	Sprlettuce.drp	Flettuce.drp	Fonion.drp	Msyonion.drp	Ssonion.drp
EBID_u_f	10.0	937.5	1250.0	593.75	625.0	1500.0	843.75	1031.25

	Grainsorghum.drp	Wheat.drp	Grchile.drp	Redchile.drp	Pecan.drp
EBID_u_f	50.0	115.0	13.75	4375.0	1447.68

;

- \* alfalfa yield in tons/acre
- \* barley yield in bu/acre
- \* potato yield in cwt/acre
- \* smgrainhay yield in tons/acre
- \* sorghumhay yield in tons/acre
- \* cornsilage yield in tons/acre
- \* pimacotton yield in lbs/acre
- \* uplandcotton yield in lbs/acre
- \* sprlettuce yield in cartons/acre
- \* flettuce yield in cartons/acre
- \* fonion yield in sacks/acre
- \* msyonion yield in sacks/acre
- \* ssonion yield in sacks/acre
- \* grainsorghum yield in cwt/acre
- \* wheat yield in cwt/acre
- \* grchile yield in tons/acre
- \* redchile yield in lbs/acre
- \* pecan yield in lbs/acre

\*barley and potato (SLV) yields from Mark Sperow's CSU dissertation

\*MRGCD crop yields from 2005 projected NMSU enterprise budgets

\*EBID crop yields from 2005 projected NMSU enterprise budgets

\*EBID pecan yields from James Coates and Dr. Rhonda Skaggs research on Dona Ana County pecan cost and return budgets

\*EPAG crop yields = EBID

\*MXAG crop yields = EBID

\*EBID drp yields derived from producer interviews and research

\*\*\*\*\*  
 \* AGRICULTURE MAP #3 CROP PRICES BY CROP, PRICES INDEPENDENT OF IRRIGATION TECHNOLOGY \*  
 \*\*\*\*\*

PARAMETER Price\_p(j) Crop Prices

/ Alfalfa	130.00
Barley	3.26
Potato	5.50
Smgrainhay	120.00
Sorghumhay	105.00
Cornsilage	16.00
Pimacotton	0.91
Uplandcotton	0.60
Sprlettuce	5.84
Flettuce	6.23
Fonion	6.63
Msyonion	6.38
Ssonion	6.43
Grainsorghum	3.70
Wheat	3.75
Grchile	285.00
Redchile	0.72
Pecan	2.28

/

- \* alfalfa price in \$/tons
- \* barley price in \$/bu
- \* potato price in \$/cwt
- \* smgrainhay price in \$/tons
- \* sorghumhay price in \$/tons
- \* cornsilage price in \$/tons
- \* pimacotton price in \$/lbs
- \* uplandcotton price in \$/lbs
- \* sprlettuce price in \$/cartons
- \* flettuce price in \$/cartons
- \* fonion price in \$/sacks
- \* msyonion price in \$/sacks
- \* ssonion price in \$/sacks
- \* grainsorghum price in \$/cwt
- \* wheat price in \$/cwt
- \* grchile price in \$/tons
- \* redchile price in \$/lbs
- \* pecan price in \$/lbs

\*barley and potato (SLV) prices from Mark Sperow CSU dissertation

\*MRGCD prices from 2005 projected NMSU enterprise budgets

\*EBID prices from 2005 projected NMSU enterprise budgets

\*NM pecan prices from NMDA/nass/usda

\*EPAG and MXAG uses EBID prices

\*\*\*\*\*  
 \* AGRICULTURE MAP #4 CROP INPUT COSTS BY CROP AND IRRIGATION TECHNOLOGY \*\*\*\*\*  
 \*\*\*\*\*

TABLE Inputcos\_p(ause, j,k) Input Production Costs Per Acre

	Alfalfa.fld	Barley.fld	Potato.fld	Smgrainhay.fld	Sorghumhay.fld	Cornsilage.fld		
SLV_u_f	400.00	440.10	1705.00	10000.00	10000.00	10000.00		
MRGCD_u_f	623.60	10000.00	10000.00	418.32	339.92	848.22		
EBID_u_f	649.52	10000.00	10000.00	10000.00	10000.00	10000.00		
MXAG_u_f	746.95	10000.00	10000.00	10000.00	10000.00	10000.00		
EPAG_u_f	746.95	10000.00	10000.00	10000.00	10000.00	10000.00		
+	Pimacotton.fld	Uplandcotton.fld	Sprlettuce.fld	Flettuce.fld	Fonion.fld	Msyonion.fld	Ssonion.fld	
SLV_u_f	10000.00	10000.00	10000.00	10000.00	10000.00	10000.00	10000.00	
MRGCD_u_f	10000.00	10000.00	10000.00	10000.00	10000.00	10000.00	10000.00	
EBID_u_f	768.12	828.77	3295.85	2842.40	6746.99	4197.18	5113.22	
MXAG_u_f	10000.00	10000.00	10000.00	10000.00	10000.00	10000.00	10000.00	
EPAG_u_f	768.12	828.77	3295.85	2842.40	6746.99	4197.18	5113.22	
+	Grainsorghum.fld	Wheat.fld	Grchile.fld	Redchile.fld	Pecan.fld			
SLV_u_f	10000.00	10000.00	10000.00	10000.00	10000.00			
MRGCD_u_f	10000.00	10000.00	10000.00	10000.00	10000.00			
EBID_u_f	313.33	441.61	2388.53	2050.02	1709.00			
MXAG_u_f	10000.00	10000.00	10000.00	10000.00	10000.00			
EPAG_u_f	313.33	441.61	2388.53	2050.02	1709.00			

\*\*note: starts EBID DI input production costs, from DI-derived enterprise budgets, projected

	Alfalfa.drp	Pimacotton.drp	Uplandcotton.drp	Sprlettuce.drp	Flettuce.drp	Fonion.drp	Msyonion.drp	Ssonion.drp
+								
EBID_u_f	784.94	758.63	815.45	3666.93	3278.56	7712.38	4858.34	5914.90

	Grainsorghum.drp	Wheat.drp	Grchile.drp	Redchile.drp	Pecan.drp
+					
EBID_u_f	330.62	446.78	2719.67	2260.35	2500.00

;

\* Costs are in \$/acre

\* High costs inserted for crops not grown in particular Ag nodes, ie. barley and potato for SLV only

\*SLV costs from Mark Sperow CSU dissertation

\*MRGCD costs from 2005 projected NMSU enterprise budgets

\*EBID costs from 2005 projected NMSU enterprise budgets

\*EBID pecan costs from James Coates and Dr. Rhonda Skaggs research on Dona Ana County pecan cost and return budgets, costs based on yields

\*EBID pecan drip irrigated costs = guesstimate

\*EPAG costs = EBID

\*MXAG costs = EPAG

\*Amortization Data Inputs

SCALAR Syslife / 10 /  
 Intrate / 0.075 /

PARAMETER DIsysc\_p Cost of DI system - beginning principle balance

;

DIsysc\_p = 2500 ;

\*Breakdown of amortization calculations, sinking fund computation

PARAMETER eqn1 eqn1  
 eqn2 eqn2  
 eqn3 eqn3  
 eqn4 eqn4  
 eqn5 eqn5

;

eqn1 = 1 / (1 + Intrate) ;  
 eqn2 = eqn1\*\*Syslife ;  
 eqn3 = 1 - eqn2 ;  
 eqn4 = DIsysc\_p \* Intrate ;  
 eqn5 = eqn4 / eqn3 ;

PARAMETER Annpayment\_p Annual loan payment amount

;

Annpayment\_p = (DIsysc\_p \* Intrate) / (1 - (1 / ((1 + Intrate)\*\*Syslife))) ;

\*Display Annpayment\_p ;

PARAMETER Costshare(k) Share paid by irrigator

;

f1d 0.00  
 drp 0.40

;

;

\*ie. drp = 1.00, subsidy pays for 0% of total cost of drip irrigation

\* drp = 0.00, subsidy pays for 100% of total cost of drip irrigation

\* drp = 0.10, irrigator pays for 10% of total cost of drip irrigation

PARAMETER Inputcost\_p(ause,j,k) Input production costs accounting for subsidy payment

; Inputcost\_p(ause,j,k) = Inputcos\_p(ause,j,k) + (Annpayment\_p \* Costshare(k)) ;

PARAMETER Netrev\_p (ause,j,k) Total net revenue per acre = tot rev - tot cost  
Netrev\_af\_p(ause,j,k) Total net revenue per acre foot = net rev \ water user

; Netrev\_p(ause,j,k) = Price\_p(j) \* Yield\_p(ause,j,k) - Inputcost\_p(ause,j,k) ;

\*Mexican Ag assumed to earn zero net revenue per acre (from US view) -- tracks MX water use but not benefits

Netrev\_p('MXAG\_u\_f', j,k) = 0 ;

Netrev\_af\_p(ause,j,k) = Netrev\_p(ause,j,k) / (BBa\_p(ause,j,k) + 0.001) ;

\*Display Inputcost\_p, Netrev\_p, Netrev\_af\_p ;

\*\*\*\*\*  
\* AGRICULTURE MAP #5 ACREAGE SUPPLIES BY NODE  
\*\*\*\*\*

PARAMETER landrh\_pp(use) Irrigable land in 1000s acres for each Ag area - 2005 census of Ag data

/  
SLV\_u\_f 277.28  
MRGCD\_u\_f 45.00  
EBID\_u\_f 89.33  
MXAG\_u\_f 100.00 > educated guess for MX irrigated acreage near C. Juarez, MX  
EPAG\_u\_f 37.20  
/

PARAMETER landrhs\_p(use, t) Available land at each Ag node by time period

; landrhs\_p(use, t) = landrh\_pp(use) ;

PARAMETER landuse\_p(use, j,t) Land reqts per acre

; landuse\_p(use, j,t) = 1 ;

TABLE Beta0(ause,j) Upper bound percentages

	Alfalfa	Barley	Potato	Smgrainhay	Sorghumhay	Cornsilage	Pimacotton	Uplandcotton	Sprletteuce	Fletteuce
SLV_u_f	0.60	0.28	0.28							
MRGCD_u_f	0.76			0.15	0.024	0.074				
EBID_u_f	0.24						0.14	0.15	0.01	0.01
MXAG_u_f	0.18						0.69	0.26		
EPAG_u_f	0.18						0.69	0.26		

	+Fonion	Msyonion	Ssonion	Grainsorghum	Wheat	Grchile	Redchile	Pecan
SLV_u_f								
MRGCD_u_f								
EBID_u_f	0.017	0.017	0.017	0.02	0.04	0.025	0.025	0.29
MXAG_u_f								0.27
EPAG_u_f								0.27

;



\*note: upper bounds data is historical acreage of crops grown in the past, taken from the yrs 1997 and 2000  
\*data collected from nass/usda, nmda, and bureau of reclamation

\*TABLE Beta1(ause,j) Lower bound percentages of major crops in EBID

	Alfalfa	Pimacotton	Uplandcotton	Pecan
*EBID_u_f	0.14	0.025	0.09	0.15

;

PARAMETER up\_bound\_p (ause,j) Upper bound on optimized acreage by node and crop

;  
up\_bound\_p (ause,j) = landrh\_pp(ause) \* Beta0(ause,j) ;

\*PARAMETER lo\_bound\_p (ause,j) Lower bound on optimized acreage in EBID

;  
\*lo\_bound\_p (ause,j) = landrh\_pp(ause) \* Beta1(ause,j) ;

\*Display up\_bound\_p ;

PARAMETER max\_onions\_p(ause,t) maximum acreage on onions for EBID 4200K acres  
max\_chile\_p (ause,t) maximum acreage on chile for EBID 4900K acres  
max\_lettuce\_p(ause,t) maximum acreage on lettuce for EBID 1800K acres

;  
max\_onions\_p ('EBID\_u\_f',t) = 4200 ;  
max\_chile\_p ('EBID\_u\_f',t) = 4900 ;  
max\_lettuce\_p('EBID\_u\_f',t) = 1800 ;

\*\*\*\*\*  
\* ECONOMICS MAPS BEGIN \*

PARAMETER scales(use) 1000s of acres by Ag node Data source USDA 2002 Ag Census

/SLV_u_f	277.28	
ABQMI_u_f	107.00	
MRGCD_u_f	45.00	
EBID_u_f	89.33	
MXAG_u_f	100.00	> educated guess for MX irrigated acreage near C. Juarez, MX
EPMI_u_f	120.56	
EPAG_u_f	37.20	

/

PARAMETER growth(use) Annual forecast growth rate of acreage or population by node

\*change from MI numbers below to zero for base MI use without ag-MI transfers

/ SLV_u_f	0.0
ABQMI_u_f	0.0240
MRGCD_u_f	0.0
EBID_u_f	0.0
MXAG_u_f	0.0
EPMI_u_f	0.0376
EPAG_u_f	0.0

/

```
* abq 0.0240
* ep 0.0376
```

```
PARAMETER scalese(res) 1000s of households receiving environmental rec reservoir benefits in basin
```

```
/HE_res_s      1
EV_res_s       1
AB_res_s       1
CO_res_s       1
EB_res_s       1
CA_res_s       1
/
```

```
PARAMETER growthe(res)  annual forecast growth rate of population by reservoir rec area
```

```
/HE_res_s      0.00
EV_res_s       0.00
AB_res_s       0.00
CO_res_s       0.00
EB_res_s       0.00
CA_res_s       0.00
/
```

```
PARAMETER
```

```
scale(use,t)    1000s forecast acres or households by node and time
scalee(res,t)   1000s forecast households benefiting from basin reservoirs by node and time
;
```

```
scale(use,t) = ((1 + growth(use ))**(ord(t)-1)) * scales(use);
scalee(res,t) = ((1 + growthe(res))**(ord(t)-1)) * scalese(res);
```

```
* display scale, scalee;
```

```
TABLE Ben_u_p(use, *) Per acre or per hhold benefit fn params for all uses
```

```
$ontext
```

```
* zero US benefits for Mexican irrigated acreage
```

*	(\$)	(\$/ac-af)	(\$/ac-af2)	(af/ac-yr1)	(af-yr1)
	intercept	linear	quadratic	u_mb_0_a	u_mb_0
SLV_u_f	195.00	265.50	-28.94	4.58	1272
ABQMI_u_f	0.00	10843.00	-9627.00	0.56	60
MRGCD_u_f	30.00	67.00	-5.92	5.65	255
EBID_u_f	137.00	94.00	-2.50	18.80	1679
MXAG_u_f	0.00	0.00	0.00	0.00	0
EPMI_u_f	0.00	9507.00	-9392.00	0.51	61
EPAG_u_f	0.00	193.00	-21.50	4.48	166

```
;
```

```
$offtext
```

```
* zero US benefits for Mexican irrigated acreage
```

*	(\$)	(\$/ac-af)	(\$/ac-af2)	(af/ac-yr1)	(af-yr1)
	intercept	linear	quadratic	u_mb_0_a	u_mb_0
SLV_u_f	.00	100.00	.00	4.58	1272
ABQMI_u_f	.00	10843.00	-9627.00	0.56	60
MRGCD_u_f	.00	50.00	.00	5.65	255
EBID_u_f	.00	75.00	.00	18.80	1679

MXAG_u_f	.00	0.00	.00	0.00	0
EPMI_u_f	.00	9507.00	-9392.00	0.51	61
EPAG_u_f	.00	100.00	.00	4.48	166

-----  
 \* Reservoir Recreation benefits are a function of reservoir volume  
 \*  
 \* Data from Richard Cole and Frank Ward published research on Riofish, AJAE August 1997  
 \*  
 \* Form is Benefit = a + bZ + cZ\*\*2

TABLE Ben\_e\_p(res, \*) Total benefits as a fn of visits at basin reservoirs - no scaling

*	(\$)	(\$/af)	(\$/af2)	(af/yr)
	intercept	linear	quadratic	v_mb_0_a
HE_res_s	4246.98	7.359	-0.002092785	1759
EV_res_s	4246.98	7.359	-0.002092785	1759
AB_res_s	4246.98	7.359	-0.002092785	1759
CO_res_s	256.62	4.104	-0.002875613	714
EB_res_s	379.82	2.210	-0.000503852	2194
CA_res_s	379.82	2.210	-0.000503852	2194

\*\*\*\*\*

-----  
 \* Table defines energy + capital costs of at all pumping nodes  
 \* (+) means that the row adds to the column's supply costs  
 \* ( ) means that the row has no effect on the column's supply costs  
 \* Cost(use) = Cost\_u\_ep\_p \* X(pump)

TABLE Cost\_u\_ep\_p(pump, apply) Energy + capital cost per acre foot pumped at use nodes

\*\*\*\*\* Column Heads are use nodes \*\*\*\*\*

	SLV_a_f	ABQMI_a_f	MRGCD_a_f	EBID_a_f	MXAG_a_f	EPMI_a_f	EPAG_a_f
SLV_p_f	40.00						
ABQMI_p_f		100.00					
MRGCD_p_f			50.00				
EBID_p_f				50.00			
MXAG_p_f					0		
EPMI_p_f						100.00	
EPAG_p_f							50.00

-----  
 \* Table defines chemical treatment costs at all pumping nodes  
 \* (+) means that the row adds to the column's supply costs  
 \* ( ) means that the row has no effect on the column's supply costs

\* Cost(use) = Cost\_u\_et\_p \* X(pump)  
 \* alb mi pump cost based on 2.50/1000 gallons = \$815 per acre foot  
 \* elp mi pump cost based on 2.00/1000 gallons = \$652 per acre foot

TABLE Cost\_u\_tp\_p(pump, apply) Chemical treatment costs per acre foot pumped at use nodes

\*\*\*\*\* Column Heads are use nodes \*\*\*\*\*

	SLV_a_f	ABQMI_a_f	MRGCD_a_f	EBID_a_f	MXAG_a_f	EPMI_a_f	EPAG_a_f
SLV_p_f	0						
ABQMI_p_f		715.00					
MRGCD_p_f			0				
EBID_p_f				0			
MXAG_p_f					0		
EPMI_p_f						715.00	
EPAG_p_f							0

\* -----  
;
  
\* -----

\* Table defines energy + capital costs at all surface diversion (not pumping) nodes  
\* (+) means that the row adds to the column's supply costs  
\* ( ) means that the row has no effect on the column's supply costs

\* Cost(use) = Cost\_u\_ed\_p \* X(divert)

\* Cost of surface water v groundwater treatment for cities comes from  
\* Mays et al, book: Regional Water Supply Planning and Expansion Models chapter 3

TABLE Cost\_u\_ed\_p(divert, apply) Energy + capital costs per acre foot diverted at use nodes

\*alb mi diversion cost based on average price of \$2.70/1000 gall = 880/acre foot  
\*elp mi diversion cost based on average 2006 price of \$2.50/1000 gall = 815/acre foot

\*\*\*\*\* Column Heads are use nodes \*\*\*\*\*

	SLV_a_f	ABQMI_a_f	MRGCD_a_f	EBID_a_f	MXAG_a_f	EPMI_a_f	EPAG_a_f
SLV_d_f	10						
ABQMI_d_f		380					
MRGCD_d_f			10				
EBID_d_f				10			
MXAG_d_f					0		
EPMI_d_f						250.00	
EPAG_d_f							10

\* -----  
;
  
\* Table defines chemical treatment costs at all surface diversion (not pumping) nodes  
\* (+) means that the row adds to the column's supply costs  
\* ( ) means that the row has no effect on the column's supply costs

\* Cost(use) = Cost\_u\_td\_p \* X(divert)

\* Cost of surface water v groundwater treatment for cities comes from  
\* Mays et al, book: Regional Water Supply Planning and Expansion Models chapter 3

TABLE Cost\_u\_td\_p(divert, apply) Chemical treatment costs per acre foot diverted at use nodes

\*\*\*\*\* Column Heads are use nodes \*\*\*\*\*

	SLV_a_f	ABQMI_a_f	MRGCD_a_f	EBID_a_f	MXAG_a_f	EPMI_a_f	EPAG_a_f
SLV_d_f	0						
ABQMI_d_f		500.00					
MRGCD_d_f			0				
EBID_d_f				0			

```

MXAG_d_f          0
EPMI_d_f          500.00
EPAG_d_f          0
* -----
;

* Table defines pumping costs per unit depth per acre foot at all aquifer nodes -
* it's based on energy costs and pumping efficiency

* (+) means that the row adds to the column's pump costs
* ( ) means that the row has no effect on the column's supply costs

* Cost(pump) = Cost_app * D(aqf)
* Cost_app approx = 0.50. For a 100 foot deep aquifer cost is $50 per acre foot pumped

TABLE Cost_app_p(aqf, pump) GW energy + cap + op cost per extra foot depth per acre foot pumped

* 0 until (1) we replace above constant energy pump costs with these below that vary with depth
* and (2) we correctly incorporate marginal increased depths in total mc of water use

***** Column Heads are pumping nodes *****
      SLV_p_f  ABQMI_p_f  MRGCD_p_f  EBID_p_f  MXAG_p_f  EPMI_p_f  EPAG_p_f
SLV_aqf_s      0.00
ABQMI_aqf_s      0.00
MRGCD_aqf_s      0.00
EBID_aqf_s      0.00
MXAG_aqf_s      0.00
EPMI_aqf_s      0.00
EPAG_aqf_s      0.00
* -----
;

PARAMETER cost_up_p(pump, apply) Total costs per acre foot pumped at apply nodes
      cost_ud_p(divert, apply) Total costs per acre foot diverted at apply nodes
;
cost_up_p(pump, apply) = cost_u_ep_p(pump, apply) + cost_u_tp_p(pump, apply);
cost_ud_p(divert, apply) = cost_u_ed_p(divert, apply) + cost_u_td_p(divert, apply);

PARAMETER Env_cost_p(res) Environmental costs per yr pr acre foot volume managing NM State Parks

* based on $10 / FTE mgmt budget for NM State Parks. Economic Impact of NM State Parks, NMSU 2004
* (1) Mgmt Cost = 0.80 * visits, (2) visits = 22.25 * water surface area,
* (3) reservoir volume to reservoir area ratio shown in ACAP relations shown below
* result is Mgmt cost = Env_cost_p * reservoir volume. Env_cost coeff shown here
* Each extra acre foot adds less than $1 to labor costs of staffing water-based reservoirs
/
HE_res_s      0.36
EV_res_s      0.31
AB_res_s      0.50
CO_res_s      0.33
EB_res_s      0.31
CA_res_s      0.62
/

* .36, .31, .50, .33, .31, .62
* 6.26, 6.80, 5.50, 1.33, 0.31, 2.00 (gives interior soln when reservoirs start full)

```

\* -----  
 PARAMETER MAX\_PRICE\_p(use) negotiated price\af used. For 44% use\af applied - MI price = \$1\1000 gal applied

/SLV\_u\_f 1  
 ABQMI\_u\_f 740  
 MRGCD\_u\_f 1  
 EBID\_u\_f 1  
 EPMI\_u\_f 740  
 MXAG\_u\_f 1  
 EPAG\_u\_f 1  
 /

\*740/af used = 326/af (\$1/1000 gal) applied

PARAMETER MIN\_use\_p(use) min reqd need = 0.1 af\hh\year = 22 gal\person\day: MWD LA uses 80gal\p\day

\* Change this parameter from 0.1 to 0.0 for both MI nodes to simulate removal of min-use-2 tiered pricing program

/SLV\_u\_f 0.0  
 ABQMI\_u\_f 0.0  
 MRGCD\_u\_f 0.0  
 EBID\_u\_f 0.0  
 EPMI\_u\_f 0.0  
 MXAG\_u\_f 0.0  
 EPAG\_u\_f 0.0  
 /

\* 0.1 af/hh/hr = 22 gallons/person/day

\*\*\*\*\*  
 \* NEXT ARE HEADWATER INFLOWS, OTHER FLOWS, FLOW RELATIONSHIPS, AND \*  
 \* RESERVOIR STARTING VOLUMES, SIMPLE ECONOMIC VALUES PER AC FT WATER USE \*  
 \*\*\*\*\*

\* all water flows are measured in 1000s acre feet per yer  
 \* all water stocks are measured in 1000s acre feet instantaneous volume

\*\$ontext

TABLE headflows(t, inflow) Annual (hist or forecast lr ave) basin inflows -- snowpack or rain

	Rio G DNG	Conej R	CBasProj	Sangr De Chr	SJChamaIBT	Rio Chma	Jemez R	Rio Puerc	Rio Sal
	RG-DN_h_f	Conejos_h_f	CBasn_h_f	SangDC_h_f	SJCham_h_f	Chama_h_f	Jemez_h_f	Puerc_h_f	Salado_h_f
2006	659800	345760	17681	227228	106580	439000	45170	32238	40515
2007	659800	345760	17681	227228	106580	439000	45170	32238	40515
2008	659800	345760	17681	227228	106580	439000	45170	32238	40515
2009	659800	345760	17681	227228	106580	439000	45170	32238	40515
2010	659800	345760	17681	227228	106580	439000	45170	32238	40515
\$ontext									
2011	659800	345760	17681	227228	106580	439000	45170	32238	40515
2012	659800	345760	17681	227228	106580	439000	45170	32238	40515
2013	659800	345760	17681	227228	106580	439000	45170	32238	40515

```

2014  659800    345760    17681    227228    106580    439000    45170    32238    40515
2015  659800    345760    17681    227228    106580    439000    45170    32238    40515
2016  659800    345760    17681    227228    106580    439000    45170    32238    40515
2017  659800    345760    17681    227228    106580    439000    45170    32238    40515
2018  659800    345760    17681    227228    106580    439000    45170    32238    40515
2019  659800    345760    17681    227228    106580    439000    45170    32238    40515
2020  659800    345760    17681    227228    106580    439000    45170    32238    40515
2021  659800    345760    17681    227228    106580    439000    45170    32238    40515
2022  659800    345760    17681    227228    106580    439000    45170    32238    40515
2023  659800    345760    17681    227228    106580    439000    45170    32238    40515
2024  659800    345760    17681    227228    106580    439000    45170    32238    40515
2025  659800    345760    17681    227228    106580    439000    45170    32238    40515
;

```

\$offtext

**PARAMETER**

```

sourc(inflow,t)  source headwater flows swaps rows and columns (messy)
source(inflow,t) source flows with drought reductions;

sourc(inflow,t) = (1/1000) * SUM(tp $ (ORD(t) EQ (ORD(tp))), headflows(t,inflow));

```

*\* Change this parameter from 1 to 0.75 to 0.50 to simulate impacts of drought;*

```

source(inflow,t) = 1.00 * sourc(inflow,t); > 100% of normal inflows;

```

*\* display sourc, source;*

**PARAMETER** discount(t) *annual discount factor*

;

```

discount(t) = (1/(1)) ** ord(t);

```

*\*display discount;*

**SCALAR**

Mx\_treaty *US Mexico 1906 Treaty delivery reqts at US-Mexico line*

*/60/*

epsilon *Small number to avoid dividing by zero*

*/.01/*

**PARAMETER**

```

B0ar(res)  A-CAPAC intercept: Intcpt for reservoir area as linear fn of volume = 0
Blar(res)  A-CAPAC slope:      (1st order) Slope for res area = linear fn of vol = d(area)\d(vol)
;

```

```

B0ar('HE_res_s') = 0; > Heron Intercept max vol = 0.4 kk af
Blar('HE_res_s') = .01475; > Heron slope max area = 5.9 k ac

B0ar('EV_res_s') = 0; > El Vado Intercept max vol = 0.186 kk af
Blar('EV_res_s') = .01720; > El Vade slope max area = 3.200 k ac

```

```

B0ar('AB_res_s') = 0; > Abiquiu Intercept max vol = 1.535 kk af
Blar('AB_res_s') = .010119; > Abiquiu slope max ares = 15.536 k ac

B0ar('CO_res_s') = 0; > Cochiti intercept max vol = 0.719 kk af
Blar('CO_res_s') = .014798; > Cochiti slope max area = 10.636 k ac

B0ar('EB_res_s') = 0; > Elephant Butte inter max vol = 2.065 kk af
Blar('EB_res_s') = .01743; > Elephant Butte slope max area = 36.000 k ac

B0ar('CA_res_s') = 0; > Caballo intercept max vol = 0.331 kk af
Blar('CA_res_s') = .03474; > Caballo slope max area = 11.500 k ac

```

**PARAMETER** B0da(aqf) *Starting aquifer depth in feet below surface*

```

/SLV_aqf_s      200
ABQMI_aqf_s    200
MRGCD_aqf_s    200
EBID_aqf_s     200
MXAG_aqf_s     200
EPMI_aqf_s     200
EPAG_aqf_s     200
/

```

**PARAMETER** B1da(aqf) *Slope (<0) for aquifer depth as linear fn of volume = d(feet) \ d(vol)*

```

/SLV_aqf_s      -.002
ABQMI_aqf_s    -.01
MRGCD_aqf_s    -.01
EBID_aqf_s     -.01
MXAG_aqf_s     -.01
EPMI_aqf_s     -.01
EPAG_aqf_s     -.01
/

```

**PARAMETER**

z0(u) *Initial reservoir and aquifer volumes at stock nodes*  
ZMAX(u) *Maximum reservoir aquifer capacity cannot be violated in any period*  
;

```

z0('HE_res_s' ) = 109.5; > Heron starting value December 2005 (1000s af)
z0('EV_res_s' ) = 25.4; > El Vado
z0('AB_res_s' ) = 111.7; > Abiquiu
z0('CO_res_s' ) = 49.0; > Cochiti
z0('EB_res_s' ) = 191.9; > Elephant Butte
z0('CA_res_s' ) = 23.3; > Caballo

z0('SLV_aqf_s' ) = 10000; > San Luis Valley aquifer starting volume (1000s af)
z0('ABQMI_aqf_s' ) = 10000; > Albuquerque aquifer st volume
z0('MRGCD_aqf_s' ) = 10000; > MRGCD aquifer starting volume
z0('EBID_aqf_s' ) = 10000; > EBID aquifer starting volume
z0('MXAG_aqf_s' ) = 10000; > MX ag aquifer starting volume
z0('EPMI_aqf_s' ) = 10000; > EP M&I aquifer starting volume
z0('EPAG_aqf_s' ) = 10000; > TX Ag aquifer starting volume

ZMAX('HE_res_s' ) = 400; > Heron capacity

```



```
ZMAX('EV_res_s')      = 186;    > El Vado capacity
ZMAX('AB_res_s')      = 1535;   > Abiquiu capacity
ZMAX('CO_res_s')      = 719;    > Cochiti capacity
ZMAX('EB_res_s')      = 2065;   > EButte Reservoir capacity (1000s af)
ZMAX('CA_res_s')      = 331;    > Caballo capacity
```

```
ZMAX('SLV_aqf_s')     = 60000;  > upper bound on SLV valley aquifer volume
ZMAX('ABQMI_aqf_s')   = 20000;  > upper bound on Albuquerque Middle Valley Aquifer
ZMAX('MRGCD_aqf_s')   = 20000;  > upper bound on MGRCD area aquifer volume
ZMAX('EBID_aqf_s')    = 20000;  > upper bound on EBID (Mesilla) aquifer volume
ZMAX('MXAG_aqf_s')    = 20000;  > upper bound on Hueco Bolson (MX) aquifer
ZMAX('EPMI_aqf_s')    = 20000;  > upper bound on Hueco Bolson (EPTX) aquifer
ZMAX('EPAG_aqf_s')    = 20000;  > upper bound on Hueco Bolson (EPTX) aquifer
```

```
*z0(res) = zmax(res);          > starts every reservoirs at full levels
```

```
***** Section 3 *****
* The following endogenous (unknown) variables are defined *
* Their numerical values are not known til GAMS finds optimal soln *
*****
```

#### FREE VARIABLES

##### \* land variables

```
Acres_v      (use,j,k,t) Total acres production in ag areas - crop - time
```

##### \* water variables

```
X_v          (i,      t) Flow: diversion-use-return flow-seepage - etc in 1000s acre feet pr yr
X_jk_v       (i,j,k,  t) Flows: - diversion-use-return - etc by node crop and irrigation technology

Z            (u,      t) Stock: volume reservoirs and aquifers in 1000s of acre feet
Za          (res,    t) Stock: area by reservoir in 1000s acres
Zd          (aqf,    t) Stock: depth by aquifer in feet
```

##### \* Econ variables all in US \$ 1000s pr yr

```
tot_ag_ben_jk_v(use,j,k,t) Flow: gross ag benefits by use node and time WITH CROP DETAIL
Tot_ag_ben_v   (ause,t)   Flow: gross ag use benefits by use node and time based on CROP DETAIL
Total_ag_ben_v (ause,t)   Flow: gross ag use benefits by use node and time NOT BASED on CROP DETAIL
Total_mi_ben_v (muse,t)   Flow: gross M&I benefits by use node and time based on M&I benefits fn

Totben_u_v     (use,t)    Flow: gross total use realted benefits based on crop detail
Totalben_u_v   (use,t)    Flow: gross total use realted benefits excluding crop detail

Ben_u_v        (use,     t) Flow: gross use related economic benfits by node and time WITHOUT CROP DETAIL
Ben_e_v        (res,     t) Flow: gross reservoir recreation benefits by node and time

Cost_ep_v      (apply,   t) Flow: pump energy + capital use cost by node time
Cost_tp_v      (apply,   t) Flow: pump treatment use cost by node and time
Cost_ed_v      (apply,   t) Flow: surface energy + capital use cost by node and time
Cost_td_v      (apply,   t) Flow: surface treatment use cost by node and time

Cost_ap_v      (pump,    t) Flow: gw depth-dependent pump cost per foot depth per ac pumped
Cost_afp_v     (pump,    t) Flow: gw depth-dependent total pump costs
Cost_dp_v      (apply,   t) Flow: gw depth-dependent total use costs due to depth dependent pump costs
```

Cost\_a\_v (apply, t) *Flow: total gross apply costs by node and time*  
 Cost\_u\_v (use, t) *Flow: total gross use costs by node and time*  
 Cost\_e\_v (res, t) *Flow: total gross env costs by node and time*  
  
 Ave\_u\_cost\_v (use, t) *Flow: ave cost per af used by node and time - used for equity analysis*  
 Ave\_e\_cost\_v (res, t) *Flow: ave cost per af stored by node and time - used for equity analysis*  
  
 N\_use\_benefit\_v(use, t) *Flow: net use related econ benefit based on CROP DETAIL by node and time*  
 N\_use\_ben\_v (use, t) *Flow: net use-related economic benefits by node and time*  
 N\_env\_ben\_v (res, t) *Flow: net environmental benefit by node and time*  
  
 NBen\_ut\_v ( t) *Flow: net benefits based on CROP detail over nodes by time*  
 NBut\_v ( t) *Flow: net benefits over nodes by time*  
 NBe\_v ( t) *Flow: net environmental benefits over nodes by time*  
  
 NBen\_u\_v *Flow: net use benefits with crop detail over nodes and time*  
 NBu\_v *Flow: net use benefits over nodes and time*  
 NBe\_v *Flow: discounted net env benefits over nodes and time*  
  
 NBen\_v *Flow: PV net benefits over nodes and time (obj)*  
 NB\_v *Flow: PV net benefits over nodes and time (objective)*  
  
 M\_u\_ben\_v(use,t) *Flow: marginal benefits of use by node and time*  
 M\_e\_ben\_v(res,t) *Flow: marginal env benefits by node and time*  
  
 Price\_u\_v(use,t) *Flow: price of use by node and time*  
 Price\_e\_v(res,t) *Flow: price of env use by node and time*  
  
 M\_cost\_e\_v(res,t) *Flow: marginal env mgmt cost of added volume by res node and time*  
  
 M\_n\_ben\_e\_v(res,t) *Flow: marginal net env benefit by res node and time = mc - mb*

*\* Equity pricing variables*

Totrev\_v(use,t) *Flow: total revenue by node and time*  
 Totrev\_n\_v(use,t) *Flow: total revenue for basic needs' uses by node and time*  
 Netrev\_v(use,t) *Flow: Cost recovery -- total net revenue (profit) by node and time*  
 Pct\_rec\_cost\_v(use,t) *Flow: percent of costs recovered through water prices charged*  
 Equity\_v(use,t) *Flow: equity -- measured as AV Cost minus price x min required use*

*\* Hydro check in 1000s acre feet per year*

USE\_v *Total water used (ET) -- summed use over use nodes*  
 USEc\_v *Total calc water use (ET) tot hw inflows minus tot outflows*

;  
 \*\*\*\*\* Section 4 \*\*\*\*\*  
 \* The following equations state relationships among a basin's \*  
 \* hydrology, institutions, and economics \*  
 \*\*\*\*\*

**EQUATIONS**

\*\*\*\*\*  
 \* EQUATIONS NAMED

\*\*\*\*\*

\*\*\*\*\*

**\* Land Block**

\*\*\*\*\*

Land (ause, t) *Flows: Agricultural land constraint*  
Acres\_ju\_e (ause, j, t) *Flows: upper bound on land based on historical acreage*  
*\*Acres\_jl\_e (ause, j, t) Flows: lower bound on land for EBID*  
Acres\_on\_e (ause, t) *Flows: onion acreage only*  
Acres\_ch\_e (ause, t) *Flows: chile acreage only*  
Acres\_lt\_e (ause, t) *Flows: lettuce acreage only*

\*\*\*\*\*

**\* Hydrology Block**

\*\*\*\*\*

Inflows (inflow, t) *Flows: set streamflows from each source node*  
Rivers (i, t) *Flows: set hydrologic mass balance by flow node: sources = uses*  
Divs (divert, t) *Flows: set diversion levels by flow node: diversions < flows*  
Evaps (evap, t) *Flows: set reservoir evaporation losses by flow node*  
Applies (apply, t) *Flows: water applied (sources) can come from diversions or pumping*  
  
Ag\_apply\_jk(aapply, j,k,t) *Flows: water applied (uses) for crop prodxn by acreage crop and technology*  
Ag\_use\_jk (ause, j,k,t) *Flows: water used for crop prodxn by acreage crop and technology*  
Ag\_seep\_jk (aseep, j,k,t) *Flows: water seeped from crop prodxn by acreage by crop and technology*  
Ag\_ret\_jk (areturn,j,k,t) *Flows: return flow from crop prodxn by acreage by crop and technology*  
  
Ag\_apply (aapply, t) *Flows: water applied (uses) for crop prodxn summed over acreage and technology*  
Ag\_use (ause, t) *Flows: water used for crop prodxn summed over acreage and technology*  
Ag\_seep (seep, t) *Flows: water seeped in crop prodxn summed over acreage and technology*  
Ag\_ret (return, t) *Flows: water returned in crop prodxn summed over acreage and technology*  
  
MI\_Uses (muse, t) *Flows: set use levels by flow node: use = proportion of application*  
MI\_Seeps (mseep, t) *Flows: set seepage levels by flow node: seep = proportion of application*  
MI>Returns (mreturn, t) *Flows: set return flow levels by flow node: rf = prop of application*  
  
Netseeps (netseep, t) *Flows: defines net seepage levels by flow node: ns = seepage - pumping*  
Gwflows (gwflow, t) *Flows: impact on river flo by flow node from net seepage*  
Aqflows (aqflow, t) *Flows: impact on aquifer vol by flow node from net seepage*  
  
reservoirs (res, t) *Stock: reservoir acctg by stock node: reserv vol falls iff net release >0*  
aquifers (aqf, t) *Stock: aquifers at the aquifer nodes: aquifer vol rises iff net seepage >0*  
area (res, t) *Stock: area by reservoir in 1000s acres = function of volume in 1000 a-f*  
depth (aqf, t) *Stock: depth by aquifer in feet = function of volume in acre feet*

**\*\$ontext**

\*\*\*\*\*

**\* Institutions Block: rules constrain water allocations**

\*\*\*\*\*

US\_MX ( t) *(International) US: Mexico Treaty of 1906 requires 60K US deliveries to MX*  
  
Federal ( t) *(US Federal) End Species Act requires 50-100 cfs year around at SA Gauge.*  
  
CO\_NM ( t) *(Interstate) Rio Grande Compact Delivery Obligation (CO to NM) at Lobatos gauge*  
NM\_TX ( t) *(Interstate) Rio Grande Compact Delivery Obligation (NM to TX) at Elephant Butte*

D2\_NM (divert, t) (Intrastate) DII Rule NM - up to 57% of releases at Caballo go to NM Lands  
D2\_TX (divert, t) (Intrastate) DII Rule TX - up to 43% of releases at Caballo to to TX lands

\*\$offtext

\*\*\*\*\*

\* Economics Block

\*\*\*\*\*

\* split benefits by ag v mi in here

Tot\_ag\_ben\_jk\_e(ause,j,k,t) Flow: ag benefits WITH CROP AND TECHNOLOGY DETAIL  
Tot\_ag\_ben\_e (ause, t) Flow: ag benefits WITH CROP DETAIL summing over crops and technology  
Total\_ag\_ben\_e (ause, t) Flow: ag benefits without crop detail from quadratic ag benefits function  
Total\_mi\_ben\_e (muse, t) Flow: m&i benefits based on m&i quadratic benefits function

Totben\_u\_e (use,t) Total benefits by use and time BASED ON CROP DETAIL  
Totalben\_u\_e (use,t) Total benefits by use and time excluding crop detail

Ben\_e\_e (res,t) Flow: environmental benefits by reservoir node

Cost\_ep\_e(apply,t) Flow: pumping energy + capital cost by node and time  
Cost\_tp\_e(apply,t) Flow: pumping treatment cost by node and time  
Cost\_ed\_e(apply,t) Flow: surface energy + capital cost by node and time  
Cost\_td\_e(apply,t) Flow: surface treatment cost by node and time

\* start depth-dependent pump costs

Cost\_ap\_e(pump,t) Flow: pumping depth cost per acre foot pumped per unit depth  
Cost\_afp\_e(pump,t) Flow: pumping dept cost from all acre feet pumped  
Cost\_dp\_e(apply,t) Flow: apply costs based on that part of applications coming from depth dependent pumping

\* finish depth-dependent pump costs

Cost\_a\_e(apply,t) Flow: total operating cost by apply nodes by time  
Cost\_u\_e(use,t) Flow: total operating costs by use node and time  
Cost\_e\_e(res,t) Flow: total environmental cost by node and time

Ave\_u\_cost\_e(use,t) Flow: average operating costs by node and time - used for equity analysis  
Ave\_e\_cost\_e(res,t) Flow: average environmental costs by node and time - used for equaity analysis

N\_use\_benefit\_e(use,t) Flow: total net use benefits (use benefits minus op costs with CROP DETAIL by node time  
N\_use\_ben\_e(use,t) Flow: total net use benefits (use benefits minus op costs) by node and time  
N\_env\_ben\_e(res,t) Flow: total net environmental benefit by node and time

NBen\_ut\_e(t) Flow: net benefits of use and environment based on crop detail over nodes by time  
NBut\_e(t) Flow: net benefits of use and environment summed over nodes by time  
NBet\_e(t) Flow: net benefits of environment summed over nodes by time

NBen\_u\_e Flow: discounted net benfits of use based on crop detail over nodes and time  
NBut\_e Flow: discounted net benefits of use over nodes and time  
NBe\_e Flow: discounted net environmental benefit over nodes and time

NBen\_e Flow: discounted net use and env benefits over nodes based on CROP detail  
NB\_e Flow: discounted net use and environmental benefits over nodes and time (obj fn)

\*\*\*\*\*

\* Policy Analysis Block

\*\*\*\*\*

Price\_u\_e(use,t)                   Flow: Price of use by node and time = mb  
Price\_e\_e(res,t)                   Flow: price of env volume by node and time

Totrev\_e(use,t)                   Flow: total gross revenue by node and time  
Netrev\_e(use,t)                   Flow: total net revenue (profit)  
Totrev\_n\_e(use,t)                  Flow: total net revenue for needs' uses

Cost\_recov\_e(mu,t)                Flow: total revenues => total costs  
Pct\_rec\_cost\_e(use,t)              Flow: percentage of total costs recovered in revenues  
Equity\_e(use,t)                    Flow: Equity measured as ac minus price x reqd use

\*\*\*\*\*

*\* Post Optimality Block*

\*\*\*\*\*

*\* marginals used to cross check*

M\_use\_ben\_e(use,t)                Flow: marginal benefits of use by node and time  
M\_env\_ben\_e(res,t)                Flow: marginal environmental benefit by node and time  
  
Marg\_cos\_e\_e(res,t)                Flow: marginal env cost of managing for more volume by node and time  
Marg\_n\_e\_ben\_e(res,t)              Flow: marginal net env benefit by reservoir node and time (meb - mec)

*\* Accounting Block*

Balance                            Flow: total ET water use basinwide -- summed ET over use nodes  
Balance\_calc                       Flow: total inferred use basinwide -- sum inflows - sum resid riv flow

;

\*\*\*\*\*

*\* End of named equations*

\*\*\*\*\*

\*\*\*\*\*

*\* Equations defined algebraically using names from above*

\*\*\*\*\*

\*\*\*\*\*

*\* AG LAND BLOCK: upper bound on available acreage under irrigation*

\*\*\*\*\*

Land(ause, t)..           sum((j, k) \$aujk(ause,j,k), Acres\_v(ause, j,k,t) \* landuse\_p(ause,j,t)) =L= LANDRHS\_p (ause, t);  
Acres\_ju\_e(ause, j, t).. sum(       k \$aujk(ause,j,k), Acres\_v(ause, j,k,t) \* landuse\_p(ause,j,t)) =L= up\_bound\_p(ause,j);  
*\*Acres\_jl\_e(ause, j, t).. sum(       k \$aujk(ause,j,k), Acres\_v(ause, j,k,t) \* landuse\_p(ause,j,t)) =G= lo\_bound\_p(ause,j);*  
Acres\_on\_e(ause,t)..       sum((on, k) \$aujk(ause,on,k), Acres\_v('EBID\_u\_f', on,k,t)) =L= max\_onions\_p('EBID\_u\_f',t);  
Acres\_ch\_e(ause,t)..       sum((ch, k) \$aujk(ause,ch,k), Acres\_v('EBID\_u\_f', ch,k,t)) =L= max\_chile\_p ('EBID\_u\_f',t);  
Acres\_lt\_e(ause,t)..       sum((lt, k) \$aujk(ause,lt,k), Acres\_v('EBID\_u\_f', lt,k,t)) =L= max\_lettuce\_p('EBID\_u\_f',t);

\*\*\*\*\*  
 \* *HYDROLOGY BLOCK: inflows, diversions, use, outflows, reservoir and aquifer levels*  
 \*\*\*\*\*

Inflows(inflow,t).. X\_v(inflow,t) =E= source(inflow,t);

Rivers(river,t).. X\_v(river,t) =E= **sum**(inflow, Bv(inflow, river) \* X\_v(inflow, t)) +  
**sum**(riverp, Bv(riverp, river) \* X\_v(riverp, t)) +  
**sum**(divert, Bv(divert, river) \* X\_v(divert, t)) +  
**sum**(return, Bv(return, river) \* X\_v(return, t)) +  
**sum**(gwflow, Bv(gwflow, river) \* X\_v(gwflow, t)) +  
**sum**(release, Bv(release,river) \* X\_v(release,t)) ;

Divs(divert,t).. X\_v(divert,t) =L= **sum**(inflow, Bd(inflow, divert) \* X\_v(inflow, t)) +  
**sum**(river, Bd(river, divert) \* X\_v(river, t)) +  
**sum**(divertp, Bd(divertp,divert) \* X\_v(divertp,t)) +  
**sum**(return, Bd(return, divert) \* X\_v(return, t)) +  
**sum**(gwflow, Bd(gwflow, divert) \* X\_v(gwflow, t)) +  
**sum**(release, Bd(release,divert) \* X\_v(release,t)) ;

Evaps(evap,t).. X\_v(evap,t) =E= **sum**(res, Be(evap, res) \* Za(res, t)) ;

Applies(apply,t).. X\_v(apply,t) =E= **sum**(divert, Ba(divert, apply) \* X\_v(divert, t)) +  
**sum**(pump, Ba(pump, apply) \* X\_v(pump, t)) ;

\* *Ag hydrology: based on water use coefficients per acre for water applied, used, seeped(gw), and returned(sw)*  
 \* *special treatment of ag is required because different crops and technologies use different amts of water*

Ag\_apply\_jk(aapply, j,k,t)\$aajk(aapply,j,k).. X\_jk\_v(aapply, j,k,t) =E= **sum**(ause, BBa\_p(aapply, j,k)  
 \* Acres\_v(ause, j,k,t) \* ID\_ua (ause, aapply ));

Ag\_use\_jk (ause, j,k,t)\$aujk(ause,j,k).. X\_jk\_v(ause, j,k,t) =E= BBa\_p(ause, j,k)  
 \* Acres\_v(ause, j,k,t) ;

Ag\_seep\_jk (aseep, j,k,t)\$asjk(aseep,j,k).. X\_jk\_v(aseep, j,k,t) =E= **sum**(ause, BBa\_p(aseep, j,k)  
 \* Acres\_v(ause, j,k,t) \* ID\_us (ause, aseep ));

Ag\_ret\_jk (areturn,j,k,t)\$arjk(areturn,j,k).. X\_jk\_v(areturn,j,k,t) =E= **sum**(ause, BBa\_p(areturn,j,k)  
 \* Acres\_v(ause, j,k,t) \* ID\_ur (ause, areturn ));

Ag\_apply (aapply, t).. X\_v (aapply, t) =E= **sum**((j,k)\$aajk(aapply,j,k), X\_jk\_v(aapply, j,k,t));  
 Ag\_use (ause, t).. X\_v (ause, t) =E= **sum**((j,k)\$aujk(ause,j,k), X\_jk\_v(ause, j,k,t));  
 Ag\_seep (aseep, t).. X\_v (aseep, t) =E= **sum**((j,k)\$asjk(aseep,j,k), X\_jk\_v(aseep, j,k,t));  
 Ag\_ret (areturn, t).. X\_v (areturn, t) =E= **sum**((j,k)\$arjk(areturn,j,k), X\_jk\_v(areturn,j,k,t));

\* *M&I hydrology: assumes a single treatment technology for ea node's mi use*

MI\_Uses (muse, t).. X\_v(muse,t) =E= **sum**(mapply, Bu(mapply, muse ) \* X\_v(mapply, t)) ;  
 MI\_Seeps (mseep, t).. X\_v(mseep,t) =E= **sum**(mapply, Bs(mapply, mseep ) \* X\_v(mapply, t)) ;  
 MI>Returns (mreturn,t).. X\_v(mreturn,t) =E= **sum**(mapply, Br(mapply, mreturn) \* X\_v(mapply, t)) ;

\* *Hydrology that applies to both ag and mi nodes*

Netseeps(netseep,t).. X\_v(netseep,t)=E= **sum**(seep, Bn(seep, netseep) \* X\_v(seep, t)) +  
**sum**(pump, Bn(pump, netseep) \* X\_v(pump, t)) ;

```
Gwflows(gwflow,t).. X_v(gwflow,t) =E= sum(netseep, Bg(netseep, gwflow)* X_v(netseep,t));
Aqflows(aqflow,t).. X_v(aqflow,t) =E= sum(netseep, Bq(netseep, aqflow)* X_v(netseep,t));

reservoirs(res,t).. Z(res,t) =E= z0(res)$ (ORD(t) EQ 1) + Z(res,t-1)
+ sum(release, BLr(release, res)* X_v(release, t))
+ sum(evap, Ber(evap, res)* X_v(evap, t));

aquifers(aqf, t).. Z(aqf,t) =E= z0(aqf)$ (ORD(t) EQ 1) + Z(aqf,t-1)
+ sum(aqflow, BQv(aqflow, aqf)* X_v(aqflow, t));
```

*\* Area of reservoirs and depth of aquifers depend on respective volumes*

```
area(res,t).. Za(res,t) =E= B0ar(res) + Blar(res) * Z(res, t);
depth(aqf,t).. Zd(aqf,t) =E= B0da(aqf) + Blda(aqf) * Z(aqf, t);
```

```
*****
* INSTITUTIONS BLOCK: Laws, compacts, treaties: rules that constrain or define use patterns
*****
*$ontext
```

*\* (International) US Mexico Treaty of 1906*

*\* Treaty text on web: <http://www.ibwc.state.gov/Files/1906Conv.pdf>  
\* Provides for 60,000 Acre Feet per year of Rio Grande delivered by US to MX at US MX Border*

```
US_MX(t).. X_v('MXAG_d_f',t) =g= Mx_treaty;
```

```
*****
* (US Federal) Endangered Species Act of 1973. Silvery Minnow needs 50 cfs year around
* Under current operation this is about 240,000 acre feet per year at the San Acacia Gauge
```

```
Federal(t).. X_v('Acacia_v_f',t) =G= 240;
```

*\* (Interstate) Rio Grande Compact of 1938 among CO, NM, AND TX*

*\* Compact text on web: <http://wrri.nmsu.edu/wrdis/compacts/Rio-Grande-Compact.pdf>*

*\* Colorado's obligation to New Mexico under the Compact, as stated in the delivery schedules, is described by a table. Here, it is approximated by a quadratic fn defining the obligation based the Rio Grande and Conejos supply indices, respectively.  
\* NM's obligations to TX exclude San Juan Chama interbasin flows from the Otowi Supply Index.  
\* SJC flows are exclusively for NM. Overdelivery is possible.*

*\* Coeffs below are based on OLS regression on flows and flows squared without intercept.  
\* Adjusted R-2s are considerably over 0.99. OSL's of t-stats are all <<.001*

```
CO_NM(t).. X_v('Lobatos_v_f',t) =G= -10
+ 0.27158 * Source('Conejos_h_f',t)
+ 0.00053735 * Source('Conejos_h_f',t) * Source('Conejos_h_f',t)
+ 0.11395 * Source('RG-DN_h_f',t)
+ 0.00032016 * Source('RG-DN_h_f',t) * Source('RG-DN_h_f',t);
```

\* New Mexico's obligation to water users below Elephant Butte Reservoir is summarized by a table,  
 \* approximated in this model by a mathematical quadratic function defining required flows delivered  
 \* to Elephant Butte Reservoir (TX) by NM, based on flows at the Otowi supply index. Overdelivery possible.

```
NM_TX(t)..      X_v('EB_g_v_f', t) - (X_v('EB_rel_f',t) + X_v('EB_evp_f',t)) =G=
0.56025 * (X_v('Otowi_v_f',t) - Source('SJCham_h_f',t))
+ 0.00010839 * (X_v('Otowi_v_f',t) - Source('SJCham_h_f',t)) * (X_v('Otowi_v_f',t) - Source('SJCham_h_f',t));
*****
```

\* (Intrastate) DII Rule: 57% max RG Project deliveries to US lands to NM Ag, 43% max to EP MI + EP AG

```
D2_NM(divert,t).. X_v('EBID_d_f', t) =L= 0.57 * (X_v('CA_g_v_f',t) - X_v('MXAG_d_f',t));
D2_TX(divert,t).. X_v('EPMI_d_f', t) + X_v('EPAG_d_f',t) =L= 0.43 * (X_v('CA_g_v_f',t) - X_v('MXAG_d_f',t));
*****
```

\*\$offtext

\*\*\*\*\*  
 \* ECONOMICS BLOCK. Everything in this block measured in monetary units  
 \*\*\*\*\*

\* Benefits by node and period

```
Tot_ag_ben_jk_e(ause,j,k,t).. tot_ag_ben_jk_v(ause,j,k,t) =E= Acres_v(ause, j,k,t)$aujk(ause,j,k) * Netrev_p(ause, j,k);
```

```
Tot_ag_ben_e (ause, t).. Tot_ag_ben_v(ause,t) =E= sum((j,k)$aujk(ause,j,k), tot_ag_ben_jk_v(ause, j,k,t));
```

```
Total_ag_ben_e (ause,t).. Total_ag_ben_v(ause, t) =E=
scale(ause,t) * ben_u_p(ause, 'intercept') +
ben_u_p(ause, 'linear') * X_v(ause,t) +
((1/scale(ause,t)) * ben_u_p(ause, 'quadratic') * X_v(ause,t) * X_v(ause,t));
```

```
Total_mi_ben_e(muse,t).. Total_mi_ben_v(muse, t) =E=
scale(muse,t) * ben_u_p(muse, 'intercept') +
ben_u_p(muse, 'linear') * X_v(muse,t) +
((1/scale(muse,t)) * ben_u_p(muse, 'quadratic') * X_v(muse,t) * X_v(muse,t));
```

```
Totben_u_e(use,t).. Totben_u_v(use,t) =E= sum (ause(use), Tot_ag_ben_v(ause, t))
+ sum (muse(use), Total_mi_ben_v(muse, t)) ;
```

```
Totalben_u_e(use,t).. Totalben_u_v(use,t) =E= sum (ause(use), Total_ag_ben_v(ause, t))
+ sum (muse(use), Total_mi_ben_v(muse, t)) ;
```

```
Ben_e_e(res,t).. Ben_e_v(res,t) =E=
scalee(res,t) * Ben_e_p(res, 'intercept') +
Ben_e_p(res, 'linear') * Z(res,t) +
(1/scalee(res,t)) * Ben_e_p(res, 'quadratic') * Z(res,t) * Z(res,t);
```

\* costs by node and period

```
Cost_ep_e(apply,t).. Cost_ep_v(apply,t) =E= sum(pump, Cost_u_ep_p(pump, apply) * X_v(pump, t));
Cost_tp_e(apply,t).. Cost_tp_v(apply,t) =E= sum(pump, Cost_u_tp_p(pump, apply) * X_v(pump, t));
Cost_ed_e(apply,t).. Cost_ed_v(apply,t) =E= sum(divert, Cost_u_ed_p(divert, apply) * X_v(divert, t));
Cost_td_e(apply,t).. Cost_td_v(apply,t) =E= sum(divert, Cost_u_td_p(divert, apply) * X_v(divert, t));
```

\* pump costs. Complicated. Increased depth increases pump costs, which reduces their use



*\* cost per added foot depth per acre foot pumped if pumping occurs*

Cost\_ap\_e(pump,t).. Cost\_ap\_v(pump,t)=E= **sum**(aqf, Cost\_app\_p(aqf, pump) \* Zd(aqf, t));

*\* cost from all acre feet pumped based on actual depth and actual pumping from that depth*

Cost\_afp\_e(pump,t).. Cost\_afp\_v(pump,t) =E= Cost\_ap\_v(pump,t) \* X\_v(pump,t);

*\* cost from total water use met by depth-dependent pumping*

Cost\_dp\_e(apply,t).. Cost\_dp\_v(apply, t) =E= **sum**(pump, ID\_pa(pump, apply) \* Cost\_afp\_v(pump,t));

*\* End of pump costs.*

*\* total costs of water uses from all sources*

Cost\_a\_e(apply,t).. Cost\_a\_v(apply,t) =E= Cost\_ep\_v(apply,t) + Cost\_tp\_v(apply,t) +  
Cost\_ed\_v(apply,t) + Cost\_td\_v(apply,t) +  
Cost\_dp\_v(apply,t) ;

Cost\_u\_e(use,t).. Cost\_u\_v(use,t) =e= **sum**(apply, ID\_au(apply, use) \* Cost\_a\_v(apply,t));

Cost\_e\_e(res,t).. Cost\_e\_v(res,t) =E= Env\_cost\_p(res) \* Z(res,t);

*\* average costs of water users from all sources - used to analyze equity*

Ave\_u\_cost\_e(use,t).. Ave\_u\_cost\_v(use,t) =E= Cost\_u\_v(use,t) / (X\_v(use,t) + epsilon);

Ave\_e\_cost\_e(res,t).. Ave\_e\_cost\_v(res,t) =E= Cost\_e\_v(res,t) / (Z(res,t) + epsilon);

\*\*\*\*\*

*\* Benefits: Both use and environmental by node and period*

N\_use\_benefit\_e(use,t).. N\_use\_benefit\_v(use, t) =E= Totben\_u\_v (use, t) - Cost\_u\_v( use,t);

N\_use\_ben\_e (use,t).. N\_use\_ben\_v (use, t) =E= Totalben\_u\_v(use, t) - Cost\_u\_v( use,t);

N\_env\_ben\_e(res,t).. N\_env\_ben\_v(res,t) =E= Ben\_e\_v(res,t) - Cost\_e\_v(res,t);

*\* Net economic benefits by time pd over nodes both for use and environmental water purposes by time*

NBen\_ut\_e(t).. NBEn\_ut\_v(t) =E= **sum**(use, N\_use\_benefit\_v(use,t));

NBut\_e(t).. NBut\_v(t) =E= **sum**(use, N\_use\_ben\_v( use,t));

NBet\_e(t).. NBet\_v(t) =E= **sum**(res, N\_env\_ben\_v(res,t));

*\* Discounted net economic and environmental benefits over time*

NBen\_u\_e.. NBEn\_u\_v =E= **sum**(t, discount(t) \* NBEn\_ut\_v(t));

NBu\_e.. NBu\_v =E= **sum**(t, discount(t) \* NBut\_v(t));

NBe\_e.. NBe\_v =E= **sum**(t, discount(t) \* NBet\_v(t));

*\* Objective fn -- sums production and environmental benefits*

NBen\_e.. NBEn\_v =E= NBEn\_u\_v + NBe\_v;

NB\_e.. NB\_v =E= NBu\_v + NBe\_v;

\*\*\*\*\*

*\* POLICY BLOCK: Various policy approaches to water pricing and allocation*

\*\*\*\*\*

*\* Revenue and costs*

*\* total revenue from poor households for needs' uses*

Totrev\_n\_e(use,t).. Totrev\_n\_v(use,t) =E= MAX\_PRICE\_p(use) \* MIN\_use\_p(use) \* Scale(use,t);  
 Totrev\_e(use,t).. Totrev\_v(use,t) =E= Price\_u\_v(use,t) \* (X\_v(use,t) - (MIN\_use\_p(use) \* Scale(use,t)))  
 + Totrev\_n\_v(use,t);  
 Netrev\_e(use,t).. Netrev\_v(use,t) =E= Totrev\_v(use,t) - Cost\_u\_v(use,t);

*\* Cost recovery criterion*

Cost\_recov\_e(mu,t).. Netrev\_v(mu,t) =G= 0;  
 Pct\_rec\_cost\_e(use,t).. Pct\_rec\_cost\_v(use,t) =E= Totrev\_v(use,t) / (Cost\_u\_v(use,t)+epsilon);

*\* Equity criterion*

Equity\_e(use,t).. Equity\_v(use,t) =E= (Ave\_u\_cost\_v(use,t) - MAX\_Price\_p(use)) \* MIN\_use\_p(use);

\*\*\*\*\* end of policy block \*\*\*\*\*

\*\*\*\*\*

*\* POST OPTIMALITY BLOCK. Nothing from here down used to find optimal solution*

\*\*\*\*\*

*\* Helps find the points where the equimarginal rule holds at both use and env nodes:*

*\* Use: MB = M Operations Cost + M Env Cost + M Res cost*  
*\* Env: MB = M Operations Cost + M Env Cost + M Res Cost*

*\* Marginal use benefits by use and time and by other subsets with time*

M\_use\_ben\_e(use,t).. M\_u\_ben\_v(use,t) =E= Ben\_u\_p( use, 'linear' )  
 + 2 \* (1/scale(use,t)) \* Ben\_u\_p( use, 'quadratic' ) \* X\_v(use,t);

*\* Marginal env benefits by reservoir and time*

M\_env\_ben\_e(res,t).. M\_e\_ben\_v(res,t) =E= Ben\_e\_p(res, 'linear')  
 + 2 \*(1/scalee(res,t)) \* Ben\_e\_p(res, 'quadratic') \* Z(res,t);

Price\_u\_e(use,t).. Price\_u\_v(use,t) =E= M\_u\_ben\_v(use,t);  
 Price\_e\_e(res,t).. Price\_e\_v(res,t) =E= M\_e\_ben\_v(res,t);

\*\*\*\*\*

*\* insert here (carefully) marginal costs from pumping-aquifer-drawdowns.*

\*

\*

\*\*\*\*\*

*\* marginal use cost is calculated AFTER optimal solution. It requires using marginals (.m)*

\*

\*

\*\*\*\*\*

*\* Marginal env (mgmt) costs by reservoir and time for volumes*

Marg\_cos\_e\_e(res,t).. M\_cost\_e\_v(res,t) =E= Env\_cost\_p(res);

\* *Marginal net env benefit at reservoir nodes*

Marg\_n\_e\_ben\_e(res,t).. M\_n\_ben\_e\_v(res,t) =E= M\_e\_ben\_v(res,t) - M\_cost\_e\_v(res,t);

\*\*\*\*\*

\* *ACCOUNTING BLOCK: tracks water flows, but not used for solution*

\*\*\*\*\*

Balance.. USE\_v =E= sum(t, sum(use, X\_v(use,t)));

Balance\_calc.. USEc\_v =E= sum(t, sum(inflow, X\_v(inflow,t))) - sum(t, X\_v('Quitman\_v\_f',t));

\*\*\*\*\*

\*\*\*\*\* *END OF ALL EQUATIONS* \*\*\*\*\*

\*\*\*\*\* *SECTION 5* \*\*\*\*\*

\* *The following section defines models.* \*

\* *Each model is defined by a set of equations used* \*

\* *for which one single variable is optimized (min or max)* \*

\*\*\*\*\*

\* *This simple prototype model uses ALL equations defined above. But larger models*

\* *may exclude some equations. For example, each of several institution could be defined*

\* *by one equation. And each of several model might conduct a single policy experiment*

\* *in which that model tries out a single institution. This would require deleting all*

\* *institutional equations except the one analyzed.*

\* *If you need to EXclude some equations, list INcluded equations where ALL appears below*

MODEL RIO\_PROTOTYPE /ALL/;

\*\*\*\*\* *Section 6* \*\*\*\*\*

\* *The following section defines all solves requested,*

\* *Each solve states a single model for which an optimum is requested.*

\*

\* *Upper, lower and fixed bounds on certain variables can also be included here*

\* *Bounding variables here gives that variable a non-zero shadow price where the optimal*

\* *solution appears at that boundary. If the bound doesn't contain the model*

\* *the variable's shadow price is zero (complementary slackness)*

\*\*\*\*\*

\* *Non-negative crop production acres*

Acres\_v.lo(use,j,k,t)=0;

\* *Non-negative flows at nodes*

X\_v.lo(inflow,t) = 0;

X\_v.lo(river,t) = 0;

X\_v.lo(divert,t) = 0;

X\_v.lo(apply,t) = 0;

X\_v.lo(use,t) = 0;

X\_v.lo(pump,t) = 0;

X\_v.lo(seep,t) = 0;

X\_v.lo(return,t) = 0;

\* *Non-negative stocks at nodes*

```
Z.lo(u,t) = 0;
Za.lo(res,t) = 0;
Zd.lo(aqf,t) = 0;
```

*\* upper and lower bounds: legal, institutional, hydrologic, or economic*

```
X_v.up('ABQMI_d_f',t) = 0; > 0 surface treatment capacity Albuquerque M&I 2006
X_v.up('MRGCD_p_f',t) = 0; > 0 pumping capacity MRGCD 2006
X_v.up('EPAG_p_f',t) = 0; > 0 pumping capacity EPaso irrigated ag 2006
X_v.up('EPMI_p_f',t) = 220; > 220K af/yr EPWU pump capacity from 74 Hueco Bolson wells
X_v.up('MXAG_p_f',t) = 0; > 0 pumping capacity from MX agriculture
X_v.up('EPMI_d_f',t) = 62; > 62K af/yr EPWU surface treatment capacity
```

*\* Sustainability terminal condition -- each water stock (reservoir, aquifer) ends with terminal volume > critical level.*

*\* Avoids depleting stocks in last period -- saves stocks for future generations*

```
Z.lo(res, t) = 0.10 * Z0( res); > each reservoir never < 10% of orig vol
Z.lo(res, tlast) = 0.50 * ZMAX(res); > term pd ea reservoir vol > 0.50 * starting value
Z.lo(aqf, tlast) = 0.50 * Z0( aqf); > term pd ea aquifer vol > 0.50 * starting value
```

```
Z.up(res, t) = ZMAX(res); > maximum reservoir capacity respected in ea period
Z.up(aqf, t) = ZMAX(aqf); > maximum aquifer capacity respected in ea period
```

*\* model run #1*

*\*OPTION NLP = conopt;*

```
SOLVE RIO_PROTOTYPE USING NLP MAXIMIZING NBen_v;
```

```
SOLVE RIO_PROTOTYPE USING NLP MAXIMIZING NB_v;
```

*\* model run #2*

```
*growth('ABQMI_u_f') = 0.0240;
*growth('EPMI_u_f') = 0.0367;
*scale(use,t) = ((1+ growth(use))**(ord(t)-1)) * scales(use);
*x.fx(ad,t) = x.l(ad,t);
*x.up(mu,t) = x.l(mu,t);
```

```
*SOLVE RIO_PROTOTYPE USING NLP MAXIMIZING NB_v;
```

```
***** Section 7 *****
* The following section displays post-optimality output *
```

**PARAMETER**

*\* table 0*

*\* calculates marginal cost from use in several steps*

```
basis_p(pump, t) Identifies interior pumping nodes not at a bound
basis_d(divert,t) Identifies interior divert nodes not at a bound
```

```

apply_marg_a(apply,t) Identifies water applied at the margin not at a bound

M_cost_a_p(apply,t) Post opt: marginal cost from appl water taken from expandable source
M_cost_ut_p(use,t) Post opt: marginal cost from used water taken from expandable source
M_cost_u_p(use) Marginal cost from water use averaged over time

M_n_ben_t_u_p(use,t) Marginal net use benefit from water use away from bounds
M_n_ben_u_p(use) Marginal net use benefit averaged over all periods

* table 1
* water quantities and some benefits

tot_applied(apply) Water applied at apply nodes
tot_divert(divert) Water diverted at divert nodes
tot_pumped(pump) Water pumped at pump nodes

eco_ben(use) Benefits at use nodes
env_ben(res) Recreation benefits at reservoir nodes

eco_sup_cost(use) Economic costs of water supply
env_cost( res) Environmental costs of env damages

* table 2
* equity pricing table

equit_v(use) Equity -- (ave cost - price) x (required water use = 0.1 af\hh\yr)
prop_rec_cos(use) Proportion of costs recovered thru water pricing

* table 3
* environmental benefits, marginal use benefits, and net benefits

Env_ben_p(res) Gross reservoir recreation benefits by node and time
Cost_p (use) Total gross economic costs by node and time
Env_cost_pp(res) Total environmental cost by node and time
Netben_p (use) Net use-related economic benefits by node and time
N_env_ben_p(res) Net environmental benefit by node and time

M_u_ben_p(use) Marginal benefits of use by node and time
M_e_ben_p(res) Marginal env benefits by node and time
M_cost_e_p(res) Marginal mgmt cost of added volume by res node and time
M_n_ben_e_p(res) Marginal net env benefit by res node and time

Netben_p (use) Net use-related economic benefits by node and time
NBu_p Net benefits over nodes by time averaged by time
NBe_p Net benefits of environmental over nodes averaged by time
NB_p Total net benefits use + environment over nodes averaged over time
;

* post optimality: parameters defined with code based on optimal solution above

* table 0

* -----
* several steps required to find marginal cost of water from expandable sources *
* -----

```

```

* Marginal use cost defined below. It's based on marginals at optimal solutions
*
* It comes by recognizing that bounded solutions will not change marginally (Kuhn Tucker: compl slackness)
* Water use expansions will not come from sources already upper or lower bounded

* 4 steps are required:

* (1): identifies interior (unbounded) optimal pumping and diversions nodes
basis_p(pump, t) = abs(X_v.m(pump, t)) lt epsilon; > un-bounded pumping
basis_d(divert,t) = abs(X_v.m(divert,t)) lt epsilon; > un-bounded diversions

* (2): finds total quantities at apply nodes met by pumping v. diversion v. a mix of both
apply_marg_a(apply,t) = sum(pump, ID_ap(apply, pump) * basis_p(pump, t) * X_v.l(pump, t)) +
                        sum(divert, ID_ad(apply, divert) * basis_d(divert,t) * X_v.l(divert,t));

* (3): finds marginal cost at apply nodes based on increases met by pumping v diversions v a mix of both
M_cost_a_p(apply,t) =
    sum(pump, Cost_up_p(pump, apply) * (X_v.l(pump, t) * basis_p(pump,t )) / (epsilon + apply_marg_a(apply,t))) +
    sum(divert, Cost_ud_p(divert, apply) * (X_v.l(divert,t) * basis_d(divert,t)) / (epsilon + apply_marg_a(apply,t))) ;

* (4): finds marginal cost at use nodes based on water used consumptively per unit applied
M_cost_ut_p(use,t) = (1/Bu_p(use)) * sum(apply, ID_au(apply, use) * M_cost_a_p(apply,t));

* averages marginal cost over all years for display
M_cost_u_p(use) = sum(t, M_cost_ut_p(use,t))/card(t);

* defines marginal net use benefits as marginal use benefits - marginal use costs;
M_n_ben_t_u_p(use,t) = M_u_ben_v.l(use,t) - M_cost_ut_p(use,t);

* averages marginal net use benefits over years for display
M_n_ben_u_p(use) = sum(t, M_n_ben_t_u_p(use,t))/(card(t) + epsilon);

* -----
* tables below are displays: they're simple averages over time of optimal solutions
* -----

* table 1

tot_applied(apply) = (epsilon + SUM(t, X_v.l(apply,t)))/card(t);
tot_divert(divert) = (epsilon + sum(t, X_v.l(divert,t)))/card(t);
tot_pumped(pump ) = (epsilon + sum(t, X_v.l(pump,t)))/card(t);

eco_ben(use) = (epsilon + sum(t, TotBen_u_v.l(use,t)))/card(t);
env_ben(res) = (epsilon + sum(t, Ben_e_v.l(res,t)))/card(t);

eco_sup_cost(use) = (epsilon + sum(t, Cost_u_v.l(use,t)))/card(t);
env_cost( res) = (epsilon + sum(t, Cost_e_v.l(res,t)))/card(t);

* table 2

equit_v(use) = (epsilon + sum(t, equity_v.l(use,t)))/card(t);
prop_rec_cos(use) = (epsilon + sum(t, pct_rec_cost_v.l(use,t)))/card(t);

* TABLE 3

```

```

Netben_p (use)      = (epsilon + sum(t,N_use_ben_v.l(use,t)))/card(t) ;
N_env_ben_p(res)   = (epsilon + sum(t,N_env_ben_v.l(res,t)))/card(t);

NBu_p              = (epsilon + sum(t,NBut_v.l(t)))/card(t);
NBe_p              = (epsilon + sum(t,NBet_v.l(t)))/card(t);
NB_p               = NBu_p + NBe_p;

M_u_ben_p(use)     = (epsilon + sum(t, M_u_ben_v.l(use,t)))/card(t);
M_e_ben_p(res)     = (epsilon + sum(t, M_e_ben_v.l(res,t)))/card(t);
M_cost_e_p(res)    = (epsilon + sum(t, M_cost_e_v.l(res,t)))/card(t);
M_n_ben_e_p(res)   = (epsilon + sum(t, M_n_ben_e_v.l(res,t)))/card(t);

DISPLAY M_cost_ut_p, M_n_ben_t_u_p;

DISPLAY tot_applied, tot_divert, tot_pumped
eco_ben,
env_ben,
eco_sup_cost, env_cost;

DISPLAY equit_v, prop_rec_cos;

DISPLAY Netben_p, N_env_ben_p
NBu_p, NBe_p, NB_p
M_u_ben_p, M_e_ben_p,
M_cost_u_p, M_cost_e_p,
M_n_ben_u_p, M_n_ben_e_p;

*****
* THE END
*****

PARAMETER Inputcostpa_p(ause,j,k,t) Input production costs per acre
;
Inputcostpa_p('EBID_u_f',j,k,t) = Inputcost_p('EBID_u_f',j,k) / landrh_pp('EBID_u_f') ;

DISPLAY Inputcostpa_p ;

```